



# JOURNAL OF AGRICULTURAL RESEARCH

VOL. XIV

WASHINGTON, D. C., SEPTEMBER 23, 1918

No. 13

## AN IMPROVED METHOD FOR RECOVERING TRYPANOSOMES FROM THE BLOOD OF RATS FOR ANTIGEN PURPOSES IN CONNECTION WITH COMPLEMENT FIXATION

By F. H. REYNOLDS and H. W. SCHOENING, *Pathological Division, Bureau of Animal Industry, United States Department of Agriculture*

### UNDESIRABLE FEATURES OF METHODS IN USE

Up to the present time the methods employed in recovering trypanosomes from the blood of artificially infected rats, though quite satisfactory, have presented some features which were somewhat detrimental to the antigen and which, if overcome, would greatly improve the same as to purity and, obviously, specificity. The question of suitable and sufficient antigens being one of vital importance, more especially in this laboratory, where many thousands of fixation tests for dourine are performed annually, it was desirable to seek an improvement.

Watson<sup>1</sup> describes a method whereby the trypanosomes might be recovered in large quantities from the blood of infected rats killed at the height of the disease. It was essentially a method of repeated centrifuging, as the erythrocytes, being of greater specific gravity than the trypanosomes, necessarily settled to the bottom. However, the method had its faults, in that many of the organisms would be drawn down with the red cells and it was necessary to sacrifice many of the trypanosomes to prevent the presence of too great a quantity of red cells in the antigen. Therefore it was necessary to eliminate the following undesirable features: First, however careful and painstaking one might be, it was quite impossible to collect all the trypanosomes present in the blood; second, it was impossible to procure a pure antigen totally devoid of rat erythrocytes which, when present in large numbers, would tend to decrease the antigenic value as well as the keeping qualities; third, it was a laborious process consuming half a day, and, as experience has shown that subjection to room temperature greatly impairs the antigenic value, the necessity for its rapid preparation and early storage on ice is quite apparent. Even then it was quite impossible to procure a pure product. There-

<sup>1</sup> WATSON, E. A. DOURINE AND THE COMPLEMENT FIXATION TEST. *In* *Parasitology*, v. 8, no. 2, p. 156-183, 1915.

fore, with a view to overcoming the several obstacles attending the preparation of the trypanosomal antigen, the following technic has been devised and has given good results.

#### TECHNIC OF NEW METHOD

Blood of infected rats is collected in a 1 per cent sodium-citrate solution in physiological salt solution in order to prevent coagulation. When all the blood has been collected, the solution is filtered through cheese-cloth to remove clots, fibrin, etc., poured into tubes, and centrifugalized for about 20 minutes at 2,100 revolutions per minute. This precipitates all the corpuscles and most of the trypanosomes, leaving an upper stratum of blood serum and citrate solution containing some of the organisms. This fluid is drawn off and again centrifugalized in order to recover any of the protozoa which may be present. To the other tubes containing the mass of corpuscles intermixed with and superimposed by trypanosomes is added sufficient distilled water to produce complete hemolysis of the rat erythrocytes, a matter of about 20 minutes, which procedure is facilitated by agitation of the mixture in a flask. This also is centrifugalized but, in this instance for about half an hour, upon the completion of which there is found at the bottom of the tubes a mass of trypanosomes with an admixture of stroma of the hemolyzed red cells, which latter, in quantity, has been found to be negligible. After discarding the supernatant fluid (hemoglobin-stained water) physiological salt solution is added and the material vigorously shaken until the mass of trypanosomes is disintegrated and evenly distributed throughout the solution. Centrifuging is again resorted to with similar results, the washed mass of trypanosomes being packed at the bottom of the tubes. The salt solution is poured off and an amount of preserving fluid (physiological salt solution and glycerin aa) equal to about twice the amount of trypanosomes added; the mixture is then agitated until a uniform suspension is acquired, when it is stored at a low temperature until used.

In order to determine whether the use of distilled water in laking the corpuscles would have any detrimental effect on the trypanosomes as regards their antigenic value, immediately following their preparation, and also after an interval of two weeks, the following procedure was employed:

Twenty white rats were inoculated subcutaneously with a suspension of *Trypanosoma equiperdum* in salt solution. At the end of the third day the blood of all the animals showed a heavy infestation with the protozoa.

The animals were then bled to death into 300 cc. of a 1 per cent sodium-citrate solution in physiological salt solution, which was then divided equally and placed into two flasks.

The trypanosomes in flask 1 were recovered by repeated centrifugalization, the machine being run a sufficient length of time to drive all the corpuscles to the bottom of the tube, but leaving a large number of trypanosomes in the supernatant fluid. This was pipetted off and the blood mixed with salt solution and again centrifugalized. By repeating this operation a number of times, as many trypanosomes as possible were obtained in suspension. This fluid was then centrifugalized, driving all the trypanosomes to the bottom of the tube. The supernatant fluid was discarded and the trypanosomes were suspended in the following solution: Glycerin 5 cc., physiological salt solution 5 cc., and labeled "Antigen I."

The trypanosomes in the second flask were recovered by breaking up the red blood cells with distilled water in the manner hereinbefore described. These trypanosomes were also suspended in 5 cc. of glycerin and 5 cc. of physiological salt solution and labeled "Antigen II."

On inspection of the two antigens it was readily seen that Antigen II contained many more trypanosomes than Antigen I and apparently no blood corpuscles, while Antigen I showed the presence of considerable blood which it had been impossible to get rid of without sacrificing many of the trypanosomes.

Both antigens were titrated for antigenic strength and anticomplementary action immediately after preparation and after being stored in the ice box for two weeks. Tests for hemolytic action were also made.

The hemolytic system employed consisted of a 3 per cent suspension of sheep red cells,  $2\frac{1}{2}$  units of hemolytic amboceptor, and  $1\frac{1}{2}$  units of complement, the latter being titrated against the amboceptor and sheep cells.

Both antigens were diluted 1 to 10 with physiological salt solution.

The test for antigenic power was made against 0.15 cc. of known positive dourine serum, which was the pooled serum from 20 horses affected with dourine.

In the test immediately after preparation the antigenic unit of Antigen I was 0.25 cc. and the anticomplementary unit was 2 cc., and was not hemolytic in five times this amount. The antigenic unit of Antigen II was 0.05 cc., the anticomplementary unit 3 cc., and was not hemolytic in five times this amount.

Two weeks after preparation the antigenic unit of Antigen I was 0.35 cc., the anticomplementary unit 1.5 cc., and the antigen showed no hemolytic action. The antigenic unit of Antigen II was 0.1 cc., the anticomplementary unit 3 cc., and it showed no hemolytic action.

The results are compared in the following table:

Antigen.	Antigenic unit.	Anticomplementary unit.	Hemolytic action.
Immediately after preparation:	Cc.	Cc.	
Antigen I. ....	0.25	2.0	None.
Antigen II. ....	.05	3.0	None.
Two weeks after preparation:			
Antigen I. ....	.35	1.5	None.
Antigen II. ....	.10	3.0	None.

The use of distilled water in laking the red blood cells had no detrimental effect on the trypanosomes with regard to their antigenic value.

#### CONCLUSIONS

In concluding, the following advantages of the new method are pointed out:

- (1) The antigen is freed of all erythrocytes.
- (2) All the trypanosomes present in the blood are recovered.
- (3) The keeping quality is improved.
- (4) The time consumed is about one and one-half hours, with practically no effort, as compared with four or five hours.
- (5) The antigenic power is increased and the anticomplementary action diminished.

## LIFE HISTORY OF PEMPHIGUS POPULI-TRANSVERSUS

By THOMAS H. JONES<sup>1</sup>

Entomological Assistant, Truck-Crop Insect Investigations, Bureau of Entomology,  
United States Department of Agriculture

WITH TECHNICAL DESCRIPTIONS BY C. P. GILLETTE, Collaborator, State Entomologist  
of Colorado

### INTRODUCTION

The first collection in Louisiana of aphids belonging to the genus *Pemphigus* from the roots of plants of the family Cruciferae seems to have been made on November 19, 1914, by Mr. E. S. Tucker, formerly Associate Entomologist of the Louisiana Experiment Stations. These specimens were taken from cabbage roots (*Brassica oleracea capitata*) at Loranger, in Tangipahoa Parish. Shortly afterwards root aphids of the same genus were noted at Baton Rouge by the writer, and material for identification was submitted to specialists acquainted with the genus. As the stages sent were not recognized as belonging to any described species, and as there were apparently no published records of any species of *Pemphigus* occurring on crucifers in the United States, studies concerning the life history, habits, and economic importance of the species were begun.

After the writer had begun the investigation Maxson (23, p. 501)<sup>2</sup> published an article on *Pemphigus betae* Doane in which he stated that—

Lice of this genus have been repeatedly taken on turnips in the south.

In a letter to the writer Mr. Maxson has given the additional information that—

the specific instances referred to were brought to my attention by Mr. F. B. Paddock, State Entomologist of Texas, he having sent material to me for identification.

<sup>1</sup> The writer wishes not only to thank Dr. Gillette for his kindness in drawing up the descriptions and supervising the preparation of illustrations, but also for suggestions regarding the biological studies, and especially for the interest which he has taken in the work. Mr. J. J. Davis, of the Bureau of Entomology, and Mr. J. R. Parker, of the Montana Agricultural Experiment Station, also have shown much interest in these studies, and the writer desires to thank them for the aid they have given. Messrs. C. E. Smith and J. L. E. Lauderdale, while members of the Bureau of Entomology, rendered valuable assistance in the studies of the life history and habits of the species.

<sup>2</sup> Reference is made by number (italic) to "Literature cited," p. 592-593.

Dr. C. P. Gillette also informed the writer that Mr. Paddock sent him material from Texas collected from turnip roots on February 13, 1914, and that the species is the same as that which attacks crucifers in Louisiana. Mr. L. C. Bragg has collected the species from watercress (*Roripa* sp.) near Fort Collins, Colo. Mr. J. J. Davis, of the Bureau of Entomology, has written that several years ago he received a species of Pemphigus collected on curly turnip (*Brassica rapa*) at Greenwood, Miss., which seems to agree with the one under discussion.

It is the purpose of this paper to present the results of investigations carried on at Baton Rouge regarding the life history and habits of the species, together with records from other points in Louisiana and from other States. These results indicate that the species of Pemphigus that feeds on the roots of crucifers is identical with the well-known *Pemphigus populi-transversus* Riley, which causes galls on the leaf petioles of some of the poplars or cottonwoods (*Populus* spp.). Fortunately, it has been possible to have these notes accompany descriptions of the various stages made by Dr. Gillette.

#### HISTORICAL REVIEW

While *Pemphigus populi-transversus* has been often referred to by entomological writers since Riley described the species in 1879 (1, p. 15-16; pl. 2, fig. 5, a-c), these references have been brief and for the most part have been limited to records of the occurrence of the species in some State or particular locality and a few words regarding bibliography and the previously recorded hosts and distribution. Such references are mentioned under "Distribution and hosts" on page 580.

In works in which insects are treated especially from an economic standpoint there are several references to the species and its gall. In 1890 Packard (3, p. 434) mentioned it in his work, "Insects injurious to forest and shade trees." Bruner (6, p. 218, fig. 57), included it in an article entitled "Insect enemies of ornamental and shade trees growing in cities and parks," which was published in 1893 in the Annual Report of the Nebraska Horticultural Society, and again mentioned it (11) in an article on aphids which appeared in The Nebraska Farmer in 1901. Lintner (9, p. 361-362) included it in his Thirteenth Report of the State Entomologist of New York, published in 1898, and stated that it had been abundant on *Populus monilifera* in Washington Park, Albany, N. Y., during 1896 and 1897. Felt (14, p. 247, 620, 635-636, pl. 11, fig. 15, 16) in his work, "Insects affecting park and woodland trees," published by the New York State Museum in 1906, also referred to its abundance in the vicinity of Albany and gave colored figures of the gall. Baldwin (21, p. 208) referred to it in the Fifth Annual Report of the State Entomologist of Indiana in 1912 and gave two original illustrations of the gall.

In lists and synopses of North American aphids the species has been mentioned by several authors. It is included in the "Host-plant list of North American Aphididae" by Williams (4, p. 6, 9), published in 1891, and in a bulletin entitled "The Aphididae of North America" by Hunter (10, p. 78), issued in 1901. In a synopsis of the genus *Pemphigus* Jackson (14, p. 182, 183, 206-208) in 1908 referred to the species somewhat in detail and stated that the life history was "very imperfectly known." Further references to its life history have been made by Davidson (16, p. 372) and Gillette and Bragg (22, p. 98). Davidson stated that the stem mothers had been observed "founding their colonies" in the vicinity of Stanford University, California, in March. Gillette and Bragg (22, p. 98) in "Notes on some Colorado aphids having alternate food plants," published in 1916, gave "winter host, *Populus* species; alternate host unknown."

#### EXPERIMENTS IN TRANSFERRING THE SPECIES FROM CRUCIFERS TO POPLAR AND FROM POPLAR TO CRUCIFERS

Soon after investigational work on this root aphid was begun, winged viviparous females (winged migrants or sexupara), collected from the soil about cruciferous roots, were sent to Dr. Gillette and to Mr. J. R. Parker. Both stated that they did not recognize the individuals as belonging to any described species of the genus *Pemphigus*, but suggested that it was possible that they might be identical with *Pemphigus populi-transversus*. In view of this, experiments were begun in an attempt to ascertain whether there was a migration of the species from crucifers to poplar at one season of the year, and a return migration at another season.

In 1916 cuttings were taken from trees of *Populus deltoides*, before the buds began to swell, and stuck in moist sand in flowerpots, which were kept in a greenhouse. When leaves began to appear on these cuttings, young individuals of the stem mother (fundatrix) were placed on them. These stem mothers had recently issued from eggs obtained in the laboratory from the true sexes, which had in turn been produced by winged females (sexupara) taken from about the roots of crucifers in the field. Swellings soon began to appear on the petioles where the stem mothers had located, and these swellings, increasing in size, gradually took on an appearance typical of the gall of *Pemphigus populi-transversus* (Pl. 82). Unfortunately the stem mothers died after the galls had reached a diameter of about  $\frac{3}{8}$  inch. Galls were again formed about stem mothers on leaf petioles of poplar in the greenhouse in the spring of 1917, and these, developing to a greater size, appeared identical to those of *P. populi-transversus*.

During the fall of 1916 winged migrants (fundatrigenia) from the galls of *Pemphigus populi-transversus* were placed under cheesecloth in a cage where turnips were growing. Later, examination of the soil showed the roots of the turnips to be heavily infested with aphids of the genus



*Pemphigus*, the infestation being similar to that which occurs on cruciferous roots under field conditions. It was evident, when the growth of the plants in this box was compared with that made by plants of the same age in an uninfested box, that they had been affected by the presence of the aphids at their roots (Pl. 84, F). In fact, the turnips died before winged migrants appeared, though probably not altogether because of the insect infestation.

Winged migrants from galls were also placed on turnips growing in pots in the greenhouse. The soil in these pots had been heated previously to a temperature sufficient to kill insect life, and the pots were covered with cloth-covered wire frames. Later, examination showed wingless individuals of *Pemphigus* sp. to be present on the roots, but again the plants died before the winged forms appeared. Before the plants died, however, a number of the wingless aphids were transferred to pots covered with cloth containing cabbage plants, the roots of which had been dipped in a mixture of water and nicotine sulphate before they were planted. On March 2 of the following year winged viviparous females, such as are found in the colonies at the roots of crucifers, were observed. No aphids appeared on the roots of control plants.

These experiments, together with others that have been carried on, indicate that the forms found on poplar and on crucifers belong to the same species. Additional proof is furnished by the fact that from poplar trees in the spring were taken winged viviparous females which agree with winged migrants found at cruciferous roots at that time of the year, and that during late summer and during the fall there were collected from the leaves of crucifers winged viviparous females which are identical with those found in galls of *P. populi-transversus*.

#### DISTRIBUTION AND HOSTS

The species has been recorded as occurring on poplar in California (16, p. 372; 19, p. 398; 20, p. 699), Texas (1, p. 15-16), Colorado (1; 7, p. 116; 22, p. 98), Kansas (12, p. 22, 23), Nebraska (18, p. 12), Missouri (1, p. 15-16), Iowa (5, p. 130), Minnesota (2, p. 20, 21), Illinois (17, p. 411), Indiana (21, p. 208), New York (9, p. 361-362; 13, p. 247, 620, 635-636; 15, p. 355), and Massachusetts (15, p. 355). Mr. Parker has collected it at Lovell, Wyoming, Dr. Gillette has specimens from Arizona, and Mr. Davis writes that he has records of its occurrence in Wisconsin, Michigan, and Ohio. Mr. H. F. Wilson has taken it in Wisconsin, and the writer has seen galls, apparently made by this species, at Agricultural College, Miss., and Jacksonville, Fla.

Four species of the genus *Populus*, *balsamifera*, *monilifera*, *trichocarpa*, and *fremontii*, have been mentioned as hosts. Britton and Brown (8, p. 491, fig. 1165; p. 493, fig. 1172) give the following distribution for *P. balsamifera*:

Newfoundland to Hudson Bay and Alaska, south to Maine, New York, Michigan, Idaho, and British Columbia.

They give *P. monilifera* as a synonym under *P. deltoides* (the name used in this article), a species which, they state, occurs from Quebec to the Northwest Territory, south to New Jersey, Florida, Colorado, and New Mexico. *P. trichocarpa* and *P. fremontii* apparently occur in the Western States.

#### FORMATION OF GALLS

In transferring the species from crucifers to sprouting cuttings of *Populus deltoides* it was found that the petioles of young leaves, just out of the bud, are apparently the only ones upon which galls begin to develop. A transverse groove first appears on the petiole where the young stem mother has located, the developing petiole gradually bending at this point. The tissue surrounding the groove, which is on the inside surface of the bent petiole, gradually enlarges until a hollow globular gall, with a transverse slit on the surface opposite the petiole, is formed around the stem mother. Galls found in the field on May 17 were roughly spherical in shape and varied in diameter from  $\frac{1}{4}$  to  $\frac{1}{2}$  inch. Just before the leaves fall from the trees some of the galls reach a greatest diameter of nearly an inch. The galls vary considerably in shape, length of the transverse slit, and development of the lips (see Pl. 83; 84, A-E). While the general color of the gall is the same as that of the petiole, a portion of the surface often has a reddish tinge.

Practically all of the galls occur somewhere on the leaf petioles, though what seem to be the galls of this species have been found on the new stem growth to which the petioles are attached. The galls usually occur singly, but as many as three have been seen in juxtaposition on one petiole.

In midsummer the gall contains, and indeed is often filled with, the comparatively large stem mother, her progeny in various stages of development, all covered with waxy secretion, together with molted skins and usually with liquid globules.

#### DATES WHEN GALLS ARE FOUND AT BATON ROUGE

In 1917, at Baton Rouge, winged viviparous females and the true sexes produced by them were taken in a small cavity on the trunk of a poplar tree as early as March 8. These winged females are identical with the winged migrants (sexupara) found about the roots of crucifers. On March 28, young stem mothers were found on leaf petioles where, to judge from the size of the galls, they had apparently been present for a few days only.

During 1916, galls were noted on May 17 which had reached a diameter of  $\frac{1}{4}$  to  $\frac{1}{2}$  inch. On November 7, when a large percentage of the leaves of the poplar had fallen, few galls remained on the trees.

## PERCENTAGE OF LEAF PETIOLES SHOWING GALLS

Leaves were gathered from a tree of *Populus deltoides* on July 26, 1916, to ascertain the percentage of leaf petioles having galls upon them, and to learn whether galls occurred in greatest numbers in any one portion of the tree as regards the height above the ground. Galls are found upon comparatively small trees and upon the largest specimens.<sup>1</sup> The tree selected was about 30 feet in height and the infestation appeared to be an average one for the particular locality. Branches were broken from it at random at various heights. The leaves were then stripped from the branches and counts made of those leaves showing galls and those not infested. It was found that there was a considerable variation in the percentage of leaves showing galls on different branches, but the distance above the ground apparently had no bearing upon the percentage of leaves infested. The results are given in Table I.

TABLE I.—Percentage of leaf petioles of *Populus deltoides* infested with *Pemphigus populi-transversus*, Baton Rouge, La., July 26, 1916

Position of branch on tree.	Number of branches examined.	Number of leaves examined.	Number of leaf petioles showing galls.	Percentage of leaf petioles showing galls.
In approximate lower third . . . . .	5	222	54	24.3
In approximate middle third . . . . .	5	441	117	26.5
In approximate upper third . . . . .	5	512	137	26.7
Total . . . . .	15	1,175	308	25.2

## DATES WHEN WINGED MIGRANTS (FUNDATRIGENIA) ARE FOUND IN GALLS

At Baton Rouge winged females begin to appear in the galls on *Populus deltoides* at a somewhat earlier date than that recorded for other localities. Riley (1, p. 15-16), in connection with his original description of the species, in which he gives Missouri, southern Texas, and Colorado as localities where this species of *Pemphigus* occurs, states that the winged females are "produced in autumn, sometimes not until the leaves have fallen." Williams (18, p. 12) mentions finding winged females, "evidently but lately transformed," in galls at Ashland, Nebr., on September 25. At Baton Rouge, in 1916, winged females were found in the galls as early as June 1, though a very small percentage of the galls contained such individuals on this date. Not until September were they present in more than 10 per cent of rather large collections examined.

<sup>1</sup> Three other species of aphids belonging to the subfamily Pemphiginae have been collected from *Populus deltoides* at Baton Rouge. Of these, *Pemphigus populicaulis* Fitch is the most common, although it is much less abundant than is *P. populi-transversus*. The two other species, found only occasionally in galls which they form on the leaves, have not been identified.

Of 200 galls collected on September 15, 17 per cent contained winged females, while they were present in all of 150 galls taken on September 28.

The number of winged females in a single gall also showed a gradual increase as the year advanced. In June usually only one was found in a gall, while on November 7 as many as 76 were present.

#### DATES WHEN WINGED MIGRANTS (FUNDATRIGENIA) LEAVE GALLS

Although winged females appeared in the galls at Baton Rouge during 1916 as early as June 1, there is no evidence to show that migration to crucifers took place until late summer. The earliest collection of winged female migrants on crucifer leaves was on August 31, and not until early October were crucifer roots found infested to any considerable extent. On October 2, 1917, during a period of clear, cool, autumn weather, the migrants from the galls were common on turnip leaves at least 500 feet from poplar trees. As many as five were found on the underside of a large leaf. The greatest migration probably occurs during October. While the writer has no definite data regarding the maximum distance they may traverse while in flight, it is probable that they, as well as the sexupara, may be carried long distances by winds.

#### NUMBER OF WINGLESS VIVIPAROUS FEMALES (VIRGOGENIA) TO WHICH MIGRANTS (FUNDATRIGENIA) GIVE BIRTH

On October 25, 1916, 25 winged females taken from galls were placed, without food, in vials and kept under observation indoors. All of these began almost immediately to give birth to young, and by October 30 all had died. The average number obtained from each individual was 26, the number ranging from 14 to 37. In one instance a female brought forth 30 young in about 24 hours.

The young viviparous females locate on the roots, feed, and when mature bring forth other wingless viviparous females. In this way the subterranean colonies become established.

#### INJURY TO AND APPEARANCE OF PLANTS INFESTED WITH SUBTERRANEAN FORMS

While severe *Pemphigus* infestation on the roots of crucifers may be indicated by the wilted condition of the leaves, a slight or moderate infestation does not usually affect, to a noticeable degree, the portions of the plant above the ground. For this reason such infestation usually goes unnoticed. In other words, this insect, while it may cause as much damage as many of those species which feed upon the leaves or other parts of the plant above the surface of the soil, does not attract as much attention as such species because it works out of the sight of the ordinary observer. Inasmuch as these aphids feed upon the roots, it is to be presumed, however, that any infestation is detrimental to the plant.

Upon examination of the soil about plants attacked by the subterranean forms colonies may be found upon any portion of the root system, but the small rootlets appear to be preferred. Where dead leaves, or other trash, occur on the surface of the soil, there is often a growth of rootlets immediately beneath, and colonies are often found in such locations.

Mr. J. L. E. Lauderdale made an interesting observation at Baton Rouge on March 19, 1917, while examining roots of *Coronopus didymus* in a field of stock beets (*Beta vulgaris*). The beets were growing on ridges where the soil was less moist and less compact than that midway between the rows. Of 25 plants of *C. didymus* growing on the ridges, 24 were infested with *Pemphigus populi-transversus*, whereas only 6 of 25 plants growing in the low ground between the rows had aphids present on their roots. This would indicate that either the compactness of the soil between the rows, or its higher moisture content, or both, was disadvantageous to the development of the aphids.

The white, flocculent material which the aphids secrete is of material aid in locating them. This secretion often occurs in considerable abundance about the colonies (Pl. 85). The light-yellow color, characteristic of the bodies of the wingless females, except in their early stages, usually makes it easy to locate them when the soil about the roots on which they are feeding is carefully examined.

#### INJURY CAUSED BY THE SUBTERRANEAN FORMS

Where the aphids occur in small or moderate numbers at the roots of plants, it is difficult to estimate the amount of damage done by them. The following extracts from correspondence received by the Louisiana Experiment Stations give information as to injury to crucifers by *Pemphigus* spp. in Louisiana. As only wingless forms were forwarded by the correspondents, it can not be stated positively that the root aphid causing the injury was *P. populi-transversus*, although that is probable.

On November 13, 1915, a correspondent living at Rhoda in St. Mary Parish wrote:

I am sending you under separate cover a cabbage plant that is badly infected with a small yellow louse, and would ask if you can recommend anything that can be done. The bug or louse is found at the root of the plant, and seems to suck the sap or eat off the roots, as the plant is badly wilted during the warm part of the day, but revives a little at night, until it finally is killed.

On November 22 of the same year a letter was received from New Iberia, Iberia Parish, in which the writer stated:

Enclosed please find stalk of cabbage with insect at the root that is destroying all my plants.

A report has also been received from Dr. C. E. Mauldin, in charge of the Iberia Live-Stock Experiment Farm of the Bureau of Animal Industry, at Jeanerette, La., that—

It has been necessary for us to abandon the planting of rape and kale at this station on account of the root-louse.

At Baton Rouge the subterranean forms apparently cause more severe injury to turnip than to any other cultivated cruciferous crop that has been under observation. Not only have they been found in greatest numbers on turnips but plants have been frequently noted which, when pulled, came up easily. The roots of these plants were mostly dead, apparently because of the attack of the aphids.

#### FOOD PLANTS OF THE SUBTERRANEAN FORMS

Wingless specimens of the genus *Pemphigus* have been taken in Louisiana from the roots of the following Cruciferae: Cabbage, turnip, mustard (*Brassica nigra*), cauliflower, and broccoli (*Brassica oleracea botrytis*), Brussels sprouts (*Brassica oleracea gemmifera*), rape (*Brassica napus*), *Coronopus didymus*, *Lepidium virginicum*, and *Roripa* sp. The last three host plants are weeds, the first being common in uncultivated fields at Baton Rouge during the winter months, when the plants are sometimes gathered and used as "greens."

Winged migrants (fundatrigenia) of the species of *Pemphigus* under consideration have been found at the roots of cabbage, turnip, Brussels sprouts, rape, *Coronopus didymus*, and *Roripa* sp. It is quite possible that further observations will disclose the fact that the species occurs also at the roots of plants not belonging to the family Cruciferae. Mr. Lauderdale has found individuals on the roots of stock beet. Adjoining infested roots of *Coronopus didymus* apparently explained their presence on the beet roots, as examination of the roots of many beets, near which no crucifers were growing, failed to disclose additional instances of infestation.

#### SPREAD OF SUBTERRANEAN FORMS

Observations made in the field and under laboratory conditions indicate that at least the smaller wingless viviparous females (virgogenia) that are present in the soil during the winter months are capable of considerable locomotion, and that when conditions become unsatisfactory, these individuals seek more suitable locations. During December they have been found in great abundance crawling over the surface of the soil and upon the plants in a field of Brussels sprouts. While carrying on some experiments in the greenhouse about a year later, individuals were found to have left flowerpots in which they were feeding on turnip roots, apparently because the turnips had begun to die as the result of being severely infested with the aphid *Myzus persicae* Sulzer. They were

especially numerous on the highest points of the cloth covering these pots, about a foot above the surface of the soil, and could be easily dislodged by slight puffs of air. Some were found under conditions which indicated that they were about 3 feet away from the nearest point where they could have originated.

As the season advances, the wingless viviparous females give birth to individuals which develop into winged viviparous females (sexupara), which later leave the soil and fly away.

#### DATES WHEN WINGED MIGRANTS (SEXUPARA) ARE FOUND IN THE SOIL

A few winged females have been found in the soil as early as December 12, and, as is the case with the winged females occurring in the galls, their number gradually increases as the season advances. During 1917, roots of *Coronopus didymus* were examined in the field from time to time with the idea of ascertaining how late in the spring the species occurs in the soil. The last winged individuals were taken on April 9. On April 16 no subterranean forms could be found, though winged migrants were alive on poplar as late as April 30.

From the field observations it appears that these winged migrants fly from the crucifers to the poplar trees during the spring, where, usually in some suitable crevice, they give birth to the true sexes. Winged females, agreeing with the winged females found about cruciferous roots, were found in such locations on poplar trees during March and April, 1917. Some observed on March 20 were dead, with true sexes and eggs located near by.

#### NUMBER OF YOUNG TO WHICH WINGED MIGRANTS (SEXUPARA) FROM CRUCIFERS GIVE BIRTH

In the laboratory, under conditions quite different from those under which the winged migrants would live in the field, the greatest number of sexed individuals to which a single aphid was observed to give birth was six. Usually this winged form brings forth all of her offspring within a short time and then dies. Those kept in the laboratory were examined daily, all of the young usually being produced from the time of one examination to the next. Individuals of both sexes have come from one winged migrant.

Examination of the abdomens of several winged migrants, collected from soil about the roots of crucifers, showed them to contain from four to seven sexed specimens, seven being the predominating number.

#### DEVELOPMENT OF THE TRUE SEXES (SEXUALES)

Eggs have been obtained in the laboratory from true sexes kept without food. The number of molts which the males and females undergo has not been ascertained, nor has it been learned when copulation takes

place. The female is larger than the male and deposits only a single egg. In a well-ventilated insectary at Baton Rouge, during 1916, eggs were first noted on March 6 in vials in which the true sexes had first appeared 12 days before.

#### OVIPOSITION

The egg is often found resting in a small amount of white, cottony material secreted by the female. The true sexes apparently do not, as a rule, move far from their places of birth, eggs being found in the field in crevices on the trunks and limbs of poplar trees where living winged migrants (sexupara) and the dead bodies of others were present. In the insectary at Baton Rouge stem mothers were first seen on March 22, 1916, in vials where eggs had first been noted on March 6, giving a period of 16 days for the incubation of the egg.

#### FORMATION OF GALL

Of necessity the young, active stem mother (fundatrix), after issuing from the egg, must make its way to the developing leaves, where it settles down on the petiole and becomes responsible for the formation of a gall. While there is no absolute proof that such is the case, it is believed from field observations that one stem mother is responsible for one gall only and that a gall is only formed when a young stem mother locates on a leaf petiole.

#### SEASONAL HISTORY OF PEMPHIGUS POPULI-TRANSVERSUS IN BRIEF

The following summary has been prepared from observations made in the field and laboratory at Baton Rouge (fig. 1). The dates when the various stages appear and migration takes place probably depend to some extent upon the weather. It would be interesting to know the seasonal history of the species in the northern portion of its range where climatic conditions, especially as regards temperature, are so different from those existing in Louisiana.

The galls begin to develop on the petioles of the young leaves of *Populus deltoides* in the spring. They increase in size during the summer and by the time the leaves fall in the autumn some have reached a diameter of nearly an inch.


Of 1,175 leaves gathered from a poplar tree on July 26, 1916, 26.2 per cent had galls on their petioles. They occur on both small and large trees.

Winged migrants (fundatrigenia) have been found in the galls as early as June 1. The percentage of galls containing winged migrants, as well as the number found in any one gall, increases as the season advances.

Winged migrants from the galls fly to various cruciferous plants. They have been found on the leaves of such plants as early as August 30 and as late as October 31.



The infestation at the roots of crucifers, which is usually made apparent by the white, cottony material which the aphids excrete, gradually becomes more severe, owing to the increase in the number of the subterranean forms. It appears that, under certain conditions, the smaller, wingless viviparous females occurring in the soil are able to migrate to a considerable distance from their place of birth and there begin new colonies.



The sexed individuals take no food and, after pairing, the female deposits a single egg.

The stem mother (fundatrix), after issuing from the egg, makes its way to the young leaves of the poplar, where it settles down on a petiole. Here a gall begins to form about it.

#### DESCRIPTIONS OF STAGES OF PEMPHIGUS POPULI-TRANSVERSUS AND ITS GALL

By C. P. GILLETTE

Examples of this species in the collection of the Colorado Experiment Station, which were collected by Mr. J. T. Monell at St. Louis, Mo., on October 2, 1907, agree in every particular with Riley's description of the winged form, except that the fifth joint of the antenna is not quite as cylindrical as the description would indicate. Specimens sent by Mr. T. H. Jones, of the Bureau of Entomology, which were collected from similar galls at Baton Rouge, agree perfectly with the Missouri specimens, and with specimens taken at many different times in Colorado. Winged and wingless "lice" taken by Mr. E. S. Tucker, then of the Bureau of Entomology, from the roots of turnips at Baton Rouge, on March 6, 1915; by Mr. F. B. Paddock, State Entomologist, on turnips, at College Station, Tex., on February 13, 1914; and by Mr. T. H. Jones on Brussels sprouts at Baton Rouge, on March 4, 1916, agree well in structural details.

The galls are widely distributed over Colorado upon the broad-leaved cottonwoods; but they are not abundant, except upon an occasional tree. It should be stated here that the sexupara of this species is separated with some difficulty from the same form of *Pemphigus betae*. In the latter species, however, the permanent sensorium of joint 5 is of the normal form and never broad and irregular, inclosing chitinous islands, as in *populi-transversus*, and the spur is distinctly longer.

**FUNDATRIX, FIRST INSTAR.**—Described from specimens reared by Mr. T. H. Jones at Baton Rouge, La., and taken from the galls on March 3, 1916.

Ground color pale yellow tinged with green; head black; wax plates dusky; length, 0.60 mm.; width, 0.23 mm.; antenna, 0.45 mm., 4-jointed, joints 1, 2, and 3 subequal in length, joint 4 with spur, almost as long as 2 and 3 combined and very stout, and with several transverse rows of small chitinous points; all femora stout, the greatest width of the hind femur nearly equalling one-half its length; six longitudinal rows of wax plates upon the dorsum of segments 1 to 6 of the abdomen, and four rows of larger plates upon the dorsum of the thorax, each plate bearing at least one short, stout hair; legs and antennae deep shining black with a few short stout gray hairs (Pl. 81, A).

**ADULT FUNDATRIX.**—Described from living specimens taken at Fort Collins, Colo., by Mr. L. C. Bragg, Assistant Entomologist of the Colorado Experiment Station, on September 22, 1916, from galls on leaves of *Populus deltoides*. The opening of the gall was a straight, transverse, or somewhat diagonal slit, passing from one-half to two-thirds of the way across the gall, but not a narrow and protruded mouthlike or liplike

opening. The galls at this date appeared to be fully grown. Besides the stem mother, there were, in each gall, a few winged "lice" quite dark in color, a good number of pupæ of varying sizes, the small ones being quite pale in color, and numerous small larvæ which were very light colored and heavily tufted with white waxy threads. The old gall mother seemed to be the sole parent of the gall colony, all of which normally acquire wings.

The stem mothers were a yellowish sordid green in color, very plump, covered with a fine white powder; head, the entire legs, including coxæ, and tips of the antennæ dusky to blackish; antenna 4-jointed and very short, not as long as hind tibia, in length approximately 0.40 mm.; length of body 2.50 to 3 mm.; joint 3 distinctly longer than joint 4 with spur, the proportion being about as 3 to 2; length of hind tibia 0.50 mm.; eyes very small. There are upon the dorsum six longitudinal rows of rather large wax plates beginning upon the mesothorax and extending to the seventh abdominal segment. Upon the prothorax and the eighth abdominal segment the number of plates is reduced to four (Pl. 81, C, H).

The writer also examined, in connection with this description, numerous specimens taken in Louisiana by Mr. T. H. Jones, in California by Messrs. E. Bethel and George P. Weldon, in Arizona by Mr. Bethel, and on the eastern and western slopes of the mountains in Colorado by Messrs. L. C. Bragg and C. P. Gillette.

It seems certain that the wingless stem mother that starts the gall of this species early in the season normally continues to feed and reproduce until the leaves mature in the fall, all of her young acquiring wings and going in search of the alternate food plants of the family Cruciferae.

**FUNDATRIGENIA MIGRANT FROM THE GALLS.**—In color and general appearance like the winged sexupara from turnips and Brussels sprouts. The specimens examined average about 0.25 mm. shorter in body length. The differences in the antennal segments are quite marked. Joint 3 has from two to six transverse sensoria, the usual number being three or four. Joint 4 is the shortest and weakest and rarely has a small sensorium. Joint 4 being somewhat smaller, and joint 5 slightly larger than in the sexupara, the contrast in size of these segments is very noticeable. The permanent sensoria on joints 5 and 6 are very large and irregular, and even may be cut into two by projecting chitinous margins. They always have upon their surfaces small chitinous pieces, one or two on joint 5 and two to four on joint 6, each bearing one or more short hairs. Upon joint 6 this large irregular sensorium may extend from the base of the spur to the middle of the segment and is nearly always very irregular in outline. The proportions of the segments are about as follows: 1, 21; 2, 30; 3, 66; 4, 31; 5, 39; 6 with spur, 68. There are many irregularities in the antennæ of this species, one of which is the frequent union of segments 3 and 4 into one (Pl. 81, J).

**WINGLESS VIVIPAROUS FEMALE.**—Described from specimens taken by Mr. T. H. Jones at Baton Rouge, La., from the roots of Brussels sprouts, on April 2, 1917.

General color sordid pale yellow, with head, antennæ, and legs dusky brown to blackish; tarsi and eyes black; length 2.50 mm.; width 1.60 mm.; antenna 0.45 mm., joints 3, and 5 plus spur, subequal; joint 4 much the shortest, being less than one-half as long as joint 3; beak barely attaining second coxæ; hind femora and tibiae

each 0.55 mm.; body and legs very free from hairs; apparently no wax glands on the body (Pl. 81, E, I).

**PUPA.**—Almost uniform pale lemon yellow with slight greenish tinge on abdomen and a shade of flesh color upon the thorax; wing pads very slightly dusky along the outer margins; head, antennæ, and all the legs dusky; eyes black.

**WINGED SEXUPARA.**—Described from living specimens taken by Mr. T. H. Jones, Baton Rouge, La., on March 21, 1917, bred from Brussels sprouts, and from preserved specimens from Mr. F. B. Paddock, College Station, Tex., which were taken on February 2, 1913; from Mr. E. S. Tucker, Baton Rouge, La., taken on March 5, 1915, on turnips; and from one specimen taken by Mr. L. C. Bragg, near Fort Collins, Colo., on watercress (*Roripa* sp.) on August 31, 1917.

Head, antenna, entire thorax above, mesothorax below, and entire legs black; wings slightly smoky, with subcostal vein black or blackish and heavy along the inner or lower margin of the stigma; abdomen sordid light greenish yellow without markings; body everywhere with a slight covering of gray powder; dorsum of abdomen covered more or less with a cottony secretion; length of body 2 mm.; wing 2.70 mm.; antenna 0.60 mm.; hind tibia 0.75 mm.; joints of antenna in following proportions: 1, 22; 2, 30; 3, 65; 4, 32; 5, 32; 6 with spur, 56, sensoria transverse, joint 3 with four to eight, usually five or six; 4 with two or three; 5 and 6, normally, with permanent sensoria only; spur near base of joint 3 distinct. Permanent sensorium on joint 5 usually very large, often inclosing one or two chitinous pieces as in fundatrigenæ. Nervures of wing dusky, the costal and subcostal being heavy and black; stigma blackish, nearly parallel sided, and about three times as long as broad (Pl. 81, D, K).

**OVIPAROUS FEMALE.**—Described from a number of specimens deposited in a cage in the laboratory by specimens sent by Mr. T. H. Jones, from Baton Rouge, La.

General color buttercup yellow, with head, antennæ, and legs whitish and very transparent; a little dusky on the vertex; eyes black; antennæ short, 0.15 mm. long, 4-jointed, joints subequal, the first, and last including the spur, longest; length of body 0.70 to 0.90 mm. (Pl. 81, G).

**EGG.**—Oblong oval, glistening, varying in color from dull white to yellow. Ten eggs, deposited in the laboratory, averaged 0.54 mm. in length, ranging from 0.48 to 0.57 mm., and 0.21 mm. in width, ranging from 0.19 to 0.24 mm.

**MALE.**—Described from specimens born along with the oviparous females.

The males differ from the females by being pale yellowish green in color, more slender in form, and a little shorter, 0.60 to 0.65 mm. long, the legs very stout (Pl. 81, F).

**GALLS.**—Riley described the gall of this species as follows:

Formed upon the petiole near the base of the leaf of *Populus monilifera* and *P. balsamifera*. An elongate-oval swelling, causing the curving and broadening of the petiole, and opening on the opposite side by a transverse slit, with a whitish, slightly thickened, and elevated margin, recalling human lips.

The writer has studied large numbers of galls of this species from Colorado and Louisiana, the latter collected by Mr. T. H. Jones. When fully grown, they vary normally from about 12 to 18 mm., extra sizes attaining 20 or even 25 mm. in their greatest diameter, which is usually

in the direction of the petiole of the leaf. The fundatrix, or stem mother, soon after hatching from the egg, locates upon the petiole of a very young opening leaf, causing it to curve and thicken, and form a transverse groove (Pl. 81, B, a, b, c) at the point of attack. The petiole continues to thicken, and the groove to deepen, forming a pit or groove which carries the "louse" with it, and the two margins or lips gradually meet, inclosing the "louse" in a spacious cavity. The mouth or slit is usually transverse, but may be turned more or less in a vertical position, and the margins may, or may not, be thickened or protruded. By the time the inmates become mature and ready to fly, the lips separate enough to allow the "lice" to pass out in search of the alternate food plants. The expanded petiole of the leaf can be easily seen extending along the convex surface of the gall opposite the mouthlike opening. (For typical forms of this gall, see Pl. 82; 84, A-E.)

The gall of this species is readily separated from the galls of *Pemphigus populicaulis*, which have a long curved opening formed by the twisting of the petiole upon itself, or from the galls of *P. populi-ramulorum*, which develop upon the side of tender growing twigs (Pl. 84, A-E).

LITERATURE CITED<sup>1</sup>

- (1) RILEY, C. V., and MONELL, J.  
1879. NOTES ON THE APHIDIDÆ OF THE UNITED STATES, WITH DESCRIPTIONS OF SPECIES OCCURRING WEST OF THE MISSISSIPPI. *In* Bul. U. S. Geol. and Geog. Survey Territories, v. 5, no. 1, p. 1-32, pl. 1-2.
- (2) OESTLUND, O. W.  
1887. SYNOPSIS OF THE APHIDIDÆ OF MINNESOTA. *Geol. and Nat. Hist. Survey Minn. Bul.* 4, 100 p. Bibliography, p. 9-12.
- (3) PACKARD, A. S.  
1890. INSECTS INJURIOUS TO FOREST AND SHADE TREES. 5th Rpt. U. S. Ent. Comm., 956 p., 306 fig., 40 pl.
- (4) WILLIAMS, T. A.  
1891. HOST-PLANT LIST OF NORTH AMERICAN APHIDIDÆ. *Univ. Nebr. Dept. Ent. Special Bul.* 1, 28 p., 1 fig.
- (5) OSBORN, Herbert.  
1892. CATALOGUE OF THE HEMIPTERA OF IOWA. *In* *Proc. Iowa Acad. Sci.*, v. 1, 1890/91, pt. 2, p. 120-131.
- (6) BRUNER, Lawrence.  
1893. INSECT ENEMIES OF ORNAMENTAL AND SHADE TREES GROWING IN CITIES AND PARKS. *In* *Ann. Rpt. Nebr. State Hort. Soc.*, 1893, p. 166-235, 72 fig.
- (7) GILLETTE, C. P., and BAKER, C. F.  
1895. A PRELIMINARY LIST OF THE HEMIPTERA OF COLORADO. *Colo. Agr. Exp. Sta. Bul.* 31 (Tech. ser. 1), 137 p., illus. Works cited, p. 5-7.
- (8) BRITTON, N. L., and BROWN, Addison.  
1896. AN ILLUSTRATED FLORA OF THE NORTHERN STATES, CANADA, AND THE BRITISH POSSESSIONS. . . . v. 1. New York.


<sup>1</sup> More recently the writer has learned that as early as 1902 Dr. F. H. Chittenden recorded the receipt of specimens of an unknown species of Pemphigus from Texas. These were sent by Mr. S. A. McHenry, of the Beeville Substation of the Texas Experiment Stations, on February 14, 1901, with the information that the species was doing injury to the roots of cabbage in the vicinity of Beeville, some of the fields being reported as totally destroyed. (CHITTENDEN, F. H. SOME INSECTS INJURIOUS TO VEGETABLE CROPS. *U. S. Dept. Agr., Div. Ent., Bul.* 33, n. 3, p. 79. 1902.)

- (9) LINTNER, J. A.  
1898. REPORT OF THE STATE ENTOMOLOGIST. *In* 51st Ann. Rpt. N. Y. State Mus., 1897, v. 1, p. 331-390, 3 fig., 2 pl.
- (10) HUNTER, W. D.  
1901. THE APHIDIDÆ OF NORTH AMERICA. Iowa Agr. Exp. Sta. Bul. 60, p. 61-138. Bibliography, p. 64-69.
- (11) BRUNER, LAWRENCE.  
1901. APHIDES, OR PLANT LICE. *In* Nebr. Farmer, v. 29, no. 26, p. 651, 657, illus.
- (12) SANBORN, C. E.  
1904. KANSAS APHIDIDÆ. *In* Kansas Univ. Sci. Bul., v. 3, no. 1, p. 3-82, pl. 1-22.
- (13) FELT, E. P.  
1905-6. INSECTS AFFECTING PARK AND WOODLAND TREES. N. Y. State Mus. Mem. 8, 2 v., 877 p., 223 fig., 70 pl.
- (14) JACKSON, C. F.  
1908. A SYNOPSIS OF THE GENUS PEMPHIGUS WITH NOTES ON THEIR ECONOMIC IMPORTANCE, LIFE HISTORY, AND GEOGRAPHICAL DISTRIBUTION. *In* Ann. Rpt. Columbus Hort. Soc., v. 22, 1907, p. 160-218, 3 fig.
- (15) GILLETTE, C. P.  
1909. PLANT LOUSE NOTES, FAMILY APHIDIDÆ. *In* Jour. Econ. Ent., v. 2, no. 5, p. 351-357, pl. 12.
- (16) DAVIDSON, W. M.  
1910. FURTHER NOTES ON THE APHIDIDÆ COLLECTED IN THE VICINITY OF STANFORD UNIVERSITY. *In* Jour. Econ. Ent., v. 3, no. 4, p. 372-381, fig. 28-29.
- (17) DAVIS, J. J.  
1910. A LIST OF THE APHIDIDÆ OF ILLINOIS, WITH NOTES ON SOME OF THE SPECIES. *In* Jour. Econ. Ent., v. 3, no. 5, p. 407-420, pl. 27-28.
- (18) WILLIAMS, T. A.  
1910. THE APHIDIDÆ OF NEBRASKA. Univ. Studies [Univ. Nebr.], v. 10, no. 2, 91 p.
- (19) DAVIDSON, W. M.  
1911. NOTES ON SOME APHIDIDÆ TAKEN IN PLACER COUNTY. *In* Pomona Coll. Jour. Ent., v. 3, no. 1, p. 398-399.
- (20) ESSIG, E. O.  
1912. APHIDIDÆ OF SOUTHERN CALIFORNIA, VIII. *In* Pomona Coll. Jour. Ent., v. 4, no. 2, p. 698-745, fig. 223-239.
- (21) BALDWIN, C. H.  
1912. FIFTH ANNUAL REPORT OF THE STATE ENTOMOLOGIST OF INDIANA, 1911-1912. 324 p., illus., 3 pl. (col.). Indianapolis.
- (22) GILLETTE, C. P., and BRAGG, L. C.  
1915. NOTES ON SOME COLORADO APHIDS HAVING ALTERNATE FOOD HABITS. *In* Jour. Econ. Ent., v. 8, no. 1, p. 97-103.
- (23) MAXSON, A. C.  
1916. SOME UNPUBLISHED NOTES ON PEMPHIGUS BETÆ DOANE. *In* Jour. Econ. Ent., v. 9, no. 5, p. 500-504.

PLATE 81

*Pemphigus populi-transversus*:

- A.—Young fundatrix, first instar.  $\times 40$ .
  - B, a, b, c.—Beginning of galls on cottonwood leaves.
  - C.—Adult fundatrix with cottony secretion removed.  $\times 13$ .
  - D.—Winged sexupara from roots of Brussels sprouts.  $\times 16$ .
  - E.—Wingless virgogene from roots of Brussels sprouts.  $\times 13$ .
  - F.—Male.  $\times 40$ .
  - G.—Oviparous female.  $\times 40$ .
  - H.—Antenna of fundatrix.  $\times 120$ .
  - I.—Antenna of wingless viviparous female from Brussels sprouts.  $\times 120$ .
  - J.—Antenna of winged fundatrigenia from gall.  $\times 120$ .
  - K.—Antenna of winged sexupara from Brussels sprouts.  $\times 120$ .
- Drawn by Miss Miriam A. Palmer.








PLATE 82

*Pemphigus populi-transversus*:


A.—Gall on poplar cutting. Cuttings were collected before the buds began to swell and were planted in sand in flowerpots kept in a greenhouse at Baton Rouge, La. Young stem mothers of *Pemphigus populi transversus*, obtained indirectly from winged female migrants from roots of crucifers, were placed on growing cuttings on March 23, 1916. The photograph, taken on May 27, shows pot containing cuttings. One developing gall can be seen on twig at upper left.

B.—Gall shown in A, enlarged to nearly natural size. Beside it can be seen the slit of a small gall which has failed to develop to any considerable extent.

PLATE 83

*Pemphigus populi-transversus:*

Variation in size of galls and location on leaf petioles of *Populus deltoides*, Baton Rouge, La., September 15, 1916. About natural size. Shows flps of galls all protruding, and in some cases thickened.



*Pemphigus populi-transversus*

*Pem. p-transversus*

PLATE 84

*Pem. p-ramulorum*





PLATE 84

A.—Gall of *Pemphigus populi-transversus*, lips not protruding.

B.—Gall of *Pemphigus populi-transversus*, lips protruding.

C.—Gall of *Pemphigus populi-ramulorum*.

D.—Beginning of gall of *Pemphigus populicaulis*.

E.—Full-grown gall of *Pemphigus populicaulis*.

All natural size. Drawn by Miriam A. Palmer.

F.—Turnip seedlings, showing injury by *Pemphigus populi-transversus*: a, infested; b, control. Turnips were planted at Baton Rouge, La., on September 16, 1916, in boxes having cloth-covered tops. Calls of *P. populi-transversus* were placed in box a on September 28. Photographed on October 13, at which time the wingless form of the aphid was abundant on roots of plants in box a. No root aphids found in box b. Note difference in growth of plants in the two boxes.

PLATE 85

*Pemphigus populi-transversus*:

F.—White cottony secretion at roots of Brussels sprouts due to presence of *Pemphigus populi-transversus*. Photographed at Baton Rouge, La., on February 10, 1916.

*Pemphigus populi-transversus*

PLATE 85



Journal of Agricultural Research

Vol. XIV, No. 13





## STEM LESIONS CAUSED BY EXCESSIVE HEAT

By CARL HARTLEY

*Pathologist, Investigations in Forest Pathology, Bureau of Plant Industry, United States Department of Agriculture*

### WHITESPOT INJURY

During work on the damping-off disease of pines (*Pinus* spp.) in 1909 the writer noticed in a nursery at Halsey, in the Nebraska sand hills, a type of disease closely corresponding to the old published descriptions of damping-off. The stems of very young seedlings developed shrunk areas at the soil surface. Commonly the entire stem was constricted by the lesion, the seedling fell over, and died. The writer has previously referred to this type of injury as "whitespot" (4, p. 5).<sup>1</sup> Close examination showed that this trouble differed in several ways from the type of damping-off which the writer and assistants have produced by inoculation with common pine-seedling parasites (*Pythium debaryanum* Hesse, *Corticium vagum* B. and C., and species of *Fusarium*). The primary whitespot lesions were in all cases limited to the stems, and usually just above the ground line. The whitespot lesion is very light in color, and this characteristic color continues to the very edge, making a sharp line of demarcation from the healthy tissue. Lesions may continue definitely limited for some days, and the upper stem and cotyledons remain turgid. In this early stage most cases of whitespot injury are easily distinguished from damping-off. Typical damping-off in porous soils is primarily a rootrot, which may attack above the ground line, but which more commonly attacks below. Damping-off lesions caused by any of the above-mentioned fungi or by *Botrytis cinerea* vary in color at different stages, gradually shading into the tissue still unaffected, and progress continuously both upward and downward.

This whitespot injury was at first supposed to be merely a special type of damping-off. Cultures made from the whitespot lesions failed to develop regularly any recognizable parasites, while most of the parallel cultures from lesions of the rootrot type yielded *Pythium debaryanum*. *Alternaria* sp. was the only fungus commonly obtained from the white spots.

Further examination showed that, in cases where whitespot lesions affected one side of the stem only, it was nearly always the south or southwest side. On seedlings which had been girdled, the lesions, if at all asymmetrical, extended higher on the south than on the north side. The south margins of the seed beds, imperfectly protected by the shade

<sup>1</sup> Reference is made by number (italic) to "Literature cited," pp. 603-604.

frames, contained more whitespot lesions than other parts of the beds. In a single bed left entirely without shade, most of the seedlings died from whitespot. These observations indicated insolation as at least a contributory cause of whitespot injury. The nursery practice involved as little watering as possible during the damping-off period. In plots given somewhat more frequent watering than the general beds, subsequent counts showed only three-fifths as much whitespot injury as elsewhere. During the three succeeding years, the nurserymen gave the seed beds much more frequent watering and more careful shading than in 1909. In careful examinations during these three years only occasional cases of whitespot were found.

The foregoing data all pointed to a physical rather than a parasitic cause for whitespot injury. Heat and light were the physical factors toward which suspicion was directed. The temperature in the surface layer of soil in the seed beds, even under the half-shade of the lath frame, was found to go as high as 52° C. The apparent preventive effect of frequent watering was believed to be due to the lowering of the soil temperature.




FIG. 1.—Lesions on seedlings of *Pinus ponderosa*: Seedlings A and D were injured by the sun's rays condensed by a lens. B was injured by a hot wire, C by an incandescent lamp, and E by the direct sun. The leaves remained turgid for 5 to 11 days after the lesions were produced. The horizontal line indicates the location of the soil surface at the time the seedlings were subjected to heat. Natural size.

In tests conducted at Washington, D. C., with seedlings of *Pinus ponderosa* typical whitespot lesions have been artificially produced. Five representative experiments are described in the following paragraphs and are illustrated in figure 1.

SEEDLING A.—The stem was subjected for less than two minutes to sunlight passed through a condensing lens at the point indicated by the constriction. The stem immediately collapsed and bent over at the point of the lesion. The soil around the root at the time was dry, and the seedling was distinctly wilted before it was heated. Later the pot was watered, and the seedling restored to a nearly vertical condition by propping. Turgor returned and was maintained for a week, but was followed by the decay of the root and wilting of the plant without extension of the original lesion (fig. 1, A).

SEEDLING B.—A heated wire held 1.5 mm. to the right of the stem for five minutes resulted in the lesion shown in figure 1, B. The plant remained turgid for 11 days and a whorl of new leaves appeared. The stem at the soil surface then decayed, apparently from a root infection, bearing a mat of pink spore masses.

SEEDLING C.—A Mazda incandescent lamp placed beside and slightly above the seedling for nearly two hours killed the tips of the cotyledons and produced a lesion just above the soil surface, without perceptible injury to intermediate tissue. In its location this lesion was thoroughly typical of those commonly found in the seed beds. Five days later fungus hyphæ appeared on the lesion, and the stem broke over at that point. The lesion then progressed both up the stem and into the root, and the seedling wilted. Spores of *Alternaria* spp. were promptly produced on placing the seedling in moist chamber (fig. 1, C).

SEEDLING D.—The stem and surrounding soil were subjected for three minutes to sunlight through a condensing lens, striking both the stem and soil surface at angles of about 45°. The resulting lesion extended about 3 mm. below the soil surface and 4 mm. above it. Mechanical support was required to keep the seedling from falling. Hyphæ soon appeared on the lesion at the soil surface, and in eight days after treatment wilting occurred, a species of *Alternaria* fruiting on the lesion in moist chamber (fig. 1, D).

SEEDLING E.—On a hot day, seedlings planted in loam in a 3 inch pot were placed immediately south of a brick wall. At night the seedling shown in figure 1, E, and another seedling were leaning slightly, but were apparently uninjured. An examination of the underground parts showed that these were distinctly shriveled from the soil surface to a point 10 mm. below; a third seedling in the pot, still erect and normal in appearance, was also constricted just below the soil surface. The plants were reported and kept under observation. The tops of all remained entirely healthy for several days. At length the lesion began to extend up the stem, and on the eighth day wilting occurred. The entire root and 10 mm. of the stem above the soil line had become involved in the original lesion. Spores of species of *Dactylosporiium*, *Alternaria*, and *Fusarium* appeared on the lesion in moist chamber.

Numerous seedlings of the same original lot were kept in the same room as the seedlings listed above during the period of the tests. None developed lesions which could have been mistaken for whitespot.

In all of the seedlings whose stems were heated directly the lesions were at first a dark grayish green, changing in 24 hours to the light color and shriveled appearance characteristic of whitespot lesions on seedlings in the nurseries. The immediate darkening is supposed to be due to the filling of the intercellular spaces with cell sap, while the ultimate light color presumably indicates the loss of liquid from both the intercellular spaces and the lumina of the cells and its replacement by air. In all cases the lesions remained definitely limited for several days, and were then extended, apparently as a result of infection by fungi not commonly capable of attacking uninjured plants. It appeared that in most cases neither the heat nor the fungi later entering the lesions stopped conduction or evolved toxins in sufficient quantity to cause the death of the leaves, as reported for another plant by Overton (11). Wilting finally occurred, it is believed, only when fungi entering at the lesion or at some point below it had penetrated the absorbing portion of the root. It may be remarked that in many cases of damping-off caused by the

ordinary seedling parasites, wilting probably occurs only after the absorbing portions of the root are invaded by the parasite.

The limitation of the whitespot lesions to the stem just above the soil surface in seedling C and in most of those observed in the nurseries, indicates that the combined radiation from the heated soil and from the sun direct ordinarily results in a temperature in the stem at that point higher than the temperature in the surface soil. In an experiment not described above, in which the source of heat was directly above the seedling, the lesion was, as might be expected, just below the soil surface rather than above it. The pot containing seedling E was also so placed that the sun's rays were more nearly perpendicular to the soil surface than in level seed beds, with the same result. It is probable that in at least some cases heat lesions will occur in the nurseries partly or entirely below the soil surface, as in seedlings D and E. It will be impossible, by any ordinary method of field observation, to distinguish from damping-off cases such as that of seedling E. Both the angle of the sun's rays, and the absorbing, conducting, and radiating capacity of the soil and of the stem will, of course, help determine whether the stem will be hottest above or just below the soil surface.

Münch (8, 9) has described the same type of injury to tree seedlings in Germany, attributing it positively to heat at the soil surface. With a thermometer having a thin, flat bulb he obtained very high temperatures in the surface soil (10). Others have also reported surface temperatures in unprotected soil from 55° to 68° C., or even higher (2, p. 55; 7, p. 13; 12; 17). Münch made an incubator test in which coniferous seedlings survived for two to three hours at temperatures not exceeding 52°, but were killed by maxima of 54° to 55° C. This seems in general to agree with the temperatures reported as being fatal to most growing plants.

Typical whitespot injury to seedlings has been found in several different States, though nowhere has it been observed to cause as heavy losses as at the Nebraska nursery, where it was first seen. Whitespot is not limited to conifers. In the vicinity of the Nebraska nursery an examination of fields of rye (*Secale cereale*) and cowpeas (*Vigna sinensis*) showed that both were affected in the seedling stage in much the same way as the pines. In the cowpeas the localization of the white, constricted lesions just above the soil surface was very marked. Plate cultures yielded no fungus suspected of parasitism. The rye seedlings were affected in the same way, though constriction was less in evidence than in the more fleshy plants. The relation between the disease and exposure to sun was very evident in the case of the rye. In the level portion of the field a moderate proportion, perhaps 5 per cent, of the shoots were affected. Where a dead furrow crossed the field from east to west this uniform distribution of disease was broken. On the wall of the dead furrow having a north exposure the disease was not noticeable.

On the other wall, exposed to the south, the percentage of affected shoots was much greater than on the level surface of the rest of the field. Wind action was not excluded as a possible cause of the cowpea lesions, but the protected location of the nursery, so far as south winds are concerned, made the evidence rather conclusive that insolation rather than wind was responsible for the lesions on the rye. Münch (8) reports whitespot on maple, vetch, and peas, and believes that in some cases germinating seeds, as well as seedlings which have already broken soil, are killed by overheated soil.

Whitespot is not always fatal, even when the lesion girdles the stem. Two seedlings of *Pinus ponderosa* which had been girdled by definite whitespot lesions, slightly shrunk but not severe enough to cause breaking over, were marked for later observation. At the end of the season the lesions had disappeared and the plants seemed in every way normal. In leaf lesions due to heat, Sorauer (15, p. 638) has after several weeks observed a regeneration of chloroplasts in slightly affected tissues.

All things considered, whitespot lesions are believed to be caused mainly by excessive heat. While light as such may possibly take part in some cases, it evidently does not enter into all cases of injury. The relative unimportance of light as distinguished from heat is indicated by the numerous lesions under slat frames, the extension of all serious lesions to the north sides of stems, and the experimental production of lesions below the soil surface (seedlings D, E, and others). The preliminary experiments here reported were mostly at excessive temperatures, and absolute proof that heat alone is the cause of the common lesions in the seed beds must await further experiments at temperatures which more commonly occur in nature.

#### BASAL LESIONS ON SEEDLINGS SEVERAL MONTHS OLD

A type of trouble which is probably related to the whitespot described in the foregoing was observed in 1915 in the seed beds of a nursery of the United States Forest Service, located at an elevation of 7,300 feet in the Wasatch Mountains, Utah. The plants affected were spruce and Douglas fir which had been raised from seed the preceding year. They had made a normal height growth during their first season and remained green throughout the winter under a heavy coating of snow which covered them for more than five months. Two or three weeks after the snow melted many of the seedlings began to turn yellow and ultimately died. Examination showed dead bark, beginning at the soil surface and extending up the stem from 3 to 9 mm. In many cases the lesion extended farther up the stem on the south than on the north side, and on some seedlings lesions were found which were entirely limited to the south side of the stem, and had started to heal over from the edges. In no case was there found any such swelling above the lesion as occurs above stem-girdle lesions on older stock. In many of the advanced cases the cortex

from the base of the stem was partly or entirely gone. Careful examination, however, indicated that even these lesions could not be attributed to any biting insect. The affected seedlings were distributed rather evenly over the beds, but in no case were any diseased plants found immediately north of posts, or on the north exposed slopes at the ends of beds, where the seedlings were somewhat protected from the sun during the hottest part of the day. Spruce, a more shade-loving tree than Douglas fir, also suffered more from the disease. The observations made indicate that the death of these seedlings was due to whiteness lesions occurring during the latter part of the preceding summer. The altitude of the nursery at first thought renders it improbable that excessive heat should have been concerned in causing the injury. While, of course, the temperature of the air at such elevations is never very high, the heat of rocks and gravel exposed to the sun at high altitudes is well known. Tubeuf (17) reports a surface soil temperature of 60° C. (140° F.) at an elevation of 10,000 feet in Yellowstone Park, 200 miles directly north of the Wasatch region.

#### BASAL STEM-GIRDLE ON OLDER STOCK

Münch and others (5, 7, p. 13; 12, 13, p. 397; 14, 15, p. 638) have further attributed to excessive surface temperatures of the soil the "*Einschnürrungskrankheit*," or stem-girdle, of older nursery stock or young forest trees of both conifers and broad-leaved trees. Dr. B. T. Galloway, of the Bureau of Plant Industry, told the writer that he had found basal lesions on young willows at Chico, Cal., which he attributed to excessive heat. In conifers this disease involves death of the base of the stem of 2-to-4-year-old seedlings and transplants. Lesions are definitely limited, and the swollen growth of the stem just above the lesion, which results from the girdling and interference with food movement, gives an appearance of constriction at the lesion itself. This disease is figured by Tubeuf (16, p. 492) and ascribed to *Pestalozzia hartigii*. Hartig (3) had originally ascribed it to the freezing of thin pools of water standing on the surface of the beds. The parasitism of *P. hartigii* has failed of confirmation (1), and Tubeuf (17) now seems to favor the view of Münch, that heat of the soil is responsible. This disease has been found at widely separated points in the United States. The writer has seen what appeared to be stem-girdle on two species of the white and three of the pitch pines, two spruces, *Abies concolor* (Gord) Parry, *Pseudotsuga taxifolia* (Poir.) Brit. *Thuja* sp., and *Juniperus* sp.<sup>1</sup> Its appearance and its ability to attack representatives of so many different genera favor a nonparasitic diagnosis. The thicker cortical tissues of the woody stems should make the cambium slower to reach maximum temperature, and prevent quite as high a temperature being reached, as in the case of the younger stems,

<sup>1</sup> Several of these observations as to coniferous species affected were communicated by Dr. J. V. Hoffman, of the Forest Service, and confirmed by the writer's examination of his specimens.

which are subject to whiteness. It is worthy of note that no cases of stem girdle have been found in four seasons' examinations of the Nebraska nursery at which whiteness has been so frequent. The heat hypothesis nevertheless seems the best explanation of stem-girdle so far offered. Tubeuf's experiments with warm water (18) are of interest in this connection as indicating that moderately high, long-continued temperatures do not necessarily kill simply by their drying effect, as has sometimes been claimed.

#### LESIONS ON UPPER PARTS OF STEMS

Older 2-needled pines and herbaceous plants as well have been observed by the writer to develop typical shrunken, definitely limited whiteness lesions on young growth of the upper parts of their stems, usually at points where an abnormal bend had made the surface nearly perpendicular to the sun's rays. Such lesions seldom girdle stems, and are rarely, if ever, of economic importance.

In the case of *Pinus strobus* the unusual amount of attention which pathologists have given it in the last two years has resulted in the finding by blister-rust scouts at five different places of yellowish lesions on young stems, sunken, and in all or nearly all cases limited to one side of the stem. Sections made by Dr. R. H. Colley, of the Bureau of Plant Industry, showed a collapsed condition of the tissues, but with absence of mycelium. Observations on these lesions in northern Wisconsin by Mr. R. G. Pierce, of the Bureau of Plant Industry, showed that practically all occur on the upper sides of bent shoots or on the west sides of vertical shoots, though a single case was found on the north side. The greater number of the affected plants were on the west sides of the nursery beds.

Most of these lesions are presumed to be due to heat. In the few cases in which soft young shoots of *Pinus banksiana* have been found girdled and bent over at the point of the lesion, mechanical bending is also suggested as a possible cause.

#### LESIONS DUE TO EXCESSIVE BENDING

Following cold weather with high winds, pine seedlings 1 to 2 weeks old have been found with white constricted basal lesions in some ways not like those produced by heat. There is reason to believe that these are due to the constant bending in high wind, at length causing the collapse of the cortical tissues without the stress being sufficient at any one time to rupture the epidermis or break the fibrovascular bundles. This is apparently an analogous case to the death of tissue between the veins of sugar maple leaves exposed to storm (6, p. 27-28). A few cases of lesions on shoots of older pines have already been mentioned as possibly the results of mechanical bending rather than of heat, and it is entirely possible that the whiteness lesions observed on cowpea seedlings should



be attributed to bending rather than to heat. Basal stem-girdle lesions found on 4-months-old wild olive seedlings (*Elaeagnus* sp.) and characterized by very slight vertical extension may be due to excessive bending. However, stem lesions caused by bending without breakage are not believed to be common enough to give rise to serious confusion with those caused by heat.

#### PREVENTIVE MEASURES

Assuming the correctness of the hypothesis that most of them are caused by heat, the logical procedure for preventing whitespot and the basal lesions on older stems is to avoid soils especially liable to overheating, and in established nurseries where trouble occurs, to artificially prevent heating. Soil with loose texture or dark surface is presumed most likely to overheat at the surface. Shading and frequent light watering have already been found helpful in preventing whitespot. Encouraging free air movement and artificially compacting the soil to increase its conducting capacity have been suggested as having prophylactic value.

#### SUMMARY

(1) Very young seedlings of conifers and certain other plants were found dying in large numbers in a Nebraska nursery from a disease which, because of its characteristic lesions, the writer has called "whitespot." The trouble has been found, though less commonly, in other localities. It is distinct from the common damping-off disease, but resembles it so closely that it is very likely to be confused with it. The lesions do not seriously interfere with the upward movement of water.

(2) The location of the whitespot lesions on the stems, their observed relation to insolation and to dry surface soil, and the production of typical lesions by artificial heating, indicate excessive heat as the cause of most of the whitespot trouble.

(3) The observation in the surface soil in the seed beds in question, and by other investigators in other places, of temperatures well over 50° C., with reported maxima as high as 68°, further substantiates the hypothesis that whitespot is due to excessive heat.

(4) Killing lesions on stems of older conifers ranging in age from several months to 4 years, are also attributed to heat. The causal importance of heat in these lesions on woody or semiwoody stems is less well established than in the case of whitespot. Further experimental work at temperatures such as actually occur in nature is necessary to settle finally the pathological importance of high soil temperature.

(5) Lesions involving young cortex and resembling those attributed to heat are probably in some cases caused by repeated bending in heavy wind without visible breakage. These are believed to be too rare to give rise to serious confusion.

## LITERATURE CITED

- (1) FISCHER, C. E. C.  
1909. NOTE ON THE BIOLOGY OF PESTALOZZIA HARTIGII, TUBEUF. *In Jour. Econ. Biol.*, v. 4, no. 3, p. 72-77, pl. 7. Literature, p. 76.
- (2) HANN, J.  
1906. *LEHRBUCH DER METEOROLOGIE*. Aufl. 2, 643 p., 89 fig., 14 pl. Leipzig.
- (3) HARTIG, Robert.  
1883. EINE NEUE ART DER FROSTBESCHÄDIGUNG IN FICHTEN- UND TANNENSAAT- UND PFLANZBEETEN. *In Allg. Forst. u. Jagdztg.*, n. F., Jahrg. 59, p. 406-409, 2 fig.
- (4) HARTLEY, Carl.  
1913. THE BLIGHTS OF CONIFEROUS NURSERY STOCK. U. S. Dept. Agr. Bull. 44, 21 p.
- (5) ———  
1916. NON-PARASITIC STEM LESIONS ON SEEDLINGS. (Abstract.) *In Phytopathology*, v. 6, no. 3, p. 308-309.
- (6) ——— and MERRILL, T. C.  
1915. STORM AND DROUGHT INJURY TO THE FOLIAGE OF ORNAMENTAL TREES. *In Phytopathology*, v. 5, no. 1, p. 20-29.
- (7) MAYR, Heinrich.  
1909. WALDBAU AUF NATURGESETZLICHER GRUNDLAGE. 568 p., 27 fig., 3 pl. (partly col.) Berlin.
- (8) MÜNCH, E.  
1913. HITZESCHÄDEN AN WALDPFLANZEN. *In Naturw. Ztschr. Forst. u. Landw.*, Jahrg. 11, Heft 12, p. 557-562, 2 fig.
- (9) ———  
1914. NOCHMAL'S HITZESCHÄDEN AN WALDPFLANZEN. *In Naturw. Ztschr. Forst. u. Landw.*, Jahrg. 12, Heft 4, p. 169-183.
- (10) ———  
1915. BEOBSACHTUNGEN ÜBER ERHITZUNG DER BOBENOBEBRFLÄCHE IN JAHRE 1914. *In Naturw. Ztschr. Forst. u. Landw.*, Jahrg. 13, Heft 6/7, p. 249-260.
- (11) OVERTON, J. B.  
1911. STUDIES ON THE RELATION OF THE LIVING CELLS TO TRANSPIRATION AND SAP-FLOW IN CYPERUS. *In Bot. Gaz.*, v. 51, no. 1, p. 28-63; no. 2, p. 102-120. Literature cited, p. 118-120.
- (12) POOL, R. J.  
1914. A STUDY OF THE VEGETATION OF THE SANDHILLS OF NEBRASKA. *In Minn. Bot. Studies*, v. 4, pt. 3, p. 189-312, illus., pl. 26-40, map. Literature cited, p. 308-312.
- (13) RAMANN, E.  
1911. *BODENKUNDE*. Aufl. 3, 699 p., 65 fig. Berlin.
- (14) SCHUSTER, Ludwig.  
1914. HITZETOD JUNGER PFLANZEN. *In Naturw. Ztschr. Forst. u. Landw.*, Jahrg. 12, Heft 8, p. 377-379, 2 fig.
- (15) SORAUER, Paul.  
1909. *HANDBUCH DER PFLANZENKRANKHEITEN*. Aufl. 3, Bd. 1. Berlin.
- (16) TUBEUF, Karl von.  
1897. DISEASES OF PLANTS INDUCED BY CRYPTOGAMIC PARASITES. English ed. by W. G. Smith, 598 p., 330 fig. London, New York.

- 
- (17) ———  
1914. HITZETOT UND EINSCHNÜRUNGSKRANKHEITEN DER PFLANZEN. In  
Naturw. Ztschr. Forst. u. Landw., Jahrg. 12, Heft 1, p. 19-36, 4 fig.
- (18) ———  
1914. ERKRANKUNGEN DURCH LUFTABSCHLUSS UND ÜBERHITZUNG. In  
Naturw. Ztschr. Forst. u. Landw., Jahrg. 12, Heft 2, p. 67-88; Heft 4,  
p. 161-169, 2 fig.

## WORK AND PARASITISM OF THE MEDITERRANEAN FRUIT FLY IN HAWAII DURING 1917

By C. E. PEMBERTON, Assistant Entomologist, and H. F. WILLARD, Fruit-Fly Quarantine Inspector, Mediterranean Fruit-Fly Investigations, Bureau of Entomology, United States Department of Agriculture

The interesting history of the many successful introductions of beneficial insects into Hawaii includes little exact data on the activities of these insects during the first few years after liberation. It is during this period that some immediate adaptations may be necessary to enable the insects to conform to a new environment, and many unexpected fluctuations may occur between the various species introduced to attack the same host before a balance is reached among them that can be expected to remain fairly constant during the years to follow. This period, then, is of much biologic interest. Apart from this, insufficient data have been published which convey accurate information concerning the work of these insects and the enormous check constantly being exerted by them over destructive pests, without which many forms of agriculture could not be conducted with profit. Aside from the aid which the entomologist can give the farmer in distributing beneficial insects over the earth beyond their natural barriers, it is his duty to obtain informing data on the kind and extent of assistance that is continually being rendered to agriculture through the work of beneficial insects, both native and introduced.

The spread and value of the parasites introduced into Hawaii to attack the Mediterranean fruit fly (*Ceratitis capitata* Wiedemann) have been watched and recorded yearly since the first liberations in 1913.<sup>1</sup> With the object of continuing this unbroken series of data and of again informing those interested of the degree of success attending these introductions, the following data are given to indicate the work of the established parasites throughout the year 1917, and the extent of fruit-fly injury caused to fruits in the Territory during that year.

During the year there was a rather heavy infestation of several varieties of fruits, some kinds being badly infested, as shown in Table I. From 913 peaches (*Amygdalus persica*) collected about Honolulu a total of 13,904 larvæ developed, or an average of 15.2 maggots to each fruit. Hardly a peach was sufficiently free from maggots to be edible. From

<sup>1</sup> BACK, E. A., and PEMBERTON, C. E. PARASITISM AMONG THE LARVÆ OF THE MEDITERRANEAN FRUIT FLY (*C. CAPITATA*) IN HAWAII IN 1914. In *Bien. Rpt. Bd. Comis. Agr. and Forestry, Hawaii, 1915-16*. D. 153-161. 1915.

BACK, E. A., and PEMBERTON, C. E. PARASITISM AMONG THE LARVÆ OF THE MEDITERRANEAN FRUIT FLY (*C. CAPITATA*) IN HAWAII IN 1915. In *Jour. Econ. Ent.*, v. 9, no. 2, p. 306-311. 1916.

PEMBERTON, C. E., and WILLARD, H. F. FRUIT-FLY PARASITISM IN HAWAII DURING 1916. In *Jour. Agr. Research*, v. 12, no. 2, p. 103-108. 1918.

17,960 kamani nuts (*Terminalia catappa*) collected during the year, 143,246 maggots were obtained, or an average of 8 to each fruit. Fortunately most other fruits were somewhat less infested than these. Though the infestation is considerable, the parasitism, as shown in Tables II and III, is very high in some cases, and during the year averaged 47.5 per cent for all fruits taken together, as determined from records on 72,139 larvæ.

TABLE I.—Extent of infestation of host fruits by larvæ of *Ceratitis capitata* in Hawaii during 1917

Host fruit.	Number of fruits collected.	Number of <i>C. capitata</i> larvæ emerging.	Average number of larvæ per fruit in 1917.	Average number of larvæ per fruit in 1916.
Indian almond ( <i>Terminalia catappa</i> )	17,960	143,246	8.0	9.5
Mango ( <i>Mangifera indica</i> )	648	5,250	8.1	1.7
Coffee ( <i>Coffea arabica</i> )	58,272	45,788	.8	.5
Strawberry guava ( <i>Psidium cattleianum</i> )	9,872	19,795	2.0	1.6
Black myrobalan ( <i>Terminalia chebula</i> )	1,767	10,411	5.9	7.0
Peach ( <i>Amygdalus persica</i> )	913	13,904	15.2	20.5
Rose-apple ( <i>Eugenia jambos</i> )	1,115	9,811	8.8	5.5
Satin-leaf ( <i>Chrysophyllum olivaceforme</i> )	6,905	23,745	3.4	2.0
French cherry ( <i>Eugenia uniflora</i> )	21,554	21,911	1.0	.8
West Indian medlar ( <i>Mimusops elengi</i> )	4,133	7,287	1.8	5.3
Yellow-wood ( <i>Ochrosia elliptica</i> )	50	1,312	23.5	3.1
Kamani ( <i>Calophyllum inophyllum</i> )	346	838	2.4	3.3
Yellow oleander ( <i>Thevetia nerifolia</i> )	2,960	16,667	5.7	3.6
Carambola ( <i>Averrhoa carambola</i> )	316	185	.6	1.3
Chinese orange ( <i>Citrus japonica</i> )	8,050	14,090	1.8	3.1
Guava ( <i>Psidium guajava</i> )	1,342	6,009	4.5	6.8
Loquat ( <i>Eriobotrya japonica</i> )	1,980	5,204	2.6	.....
Kona orange ( <i>Citrus sinensis</i> )	350	1,600	4.6	.....
Mandarin orange ( <i>Citrus nobilis deliciosa</i> )	341	732	2.2	.....
Sapodilla ( <i>Achras zapota</i> )	104	488	4.7	.....
White sapote ( <i>Casimiroa edulis</i> )	244	1,845	7.6	.....
Wampi ( <i>Clausena wampi</i> )	117	24	.2	.....

TABLE II.—Percentage of larval parasitism of *Ceratitis capitata* in Hawaii in 1917\*

Host fruit.	Month of collection.	Number of larvae emerging during first 7-6 days.	Percentage of parasitism.					Total.
			<i>Opis humilis.</i>	<i>Diachasma tryoni.</i>	<i>Diachasma fullawayi.</i>	<i>Tetrastichus gilfordianus.</i>		
	1917.							
Indian almond .....	January....	677	7.5	44.3	.....	1.3	53.6	
Do.....	February....	690	6.1	16.8	.....	1.3	24.2	
Do.....	March.....	302	10.3	22.2	.....	7.3	39.8	
Do.....	April.....	342	10.2	22.8	.....	3.5	36.5	
Do.....	May.....	1,035	11.4	34.3	0.3	.7	46.7	
Do.....	August.....	4,620	4.4	15.8	.2	3.5	23.9	
Do.....	September..	5,148	5.0	46.8	.02	3.3	55.12	

\* Most of the fruits represented in this table were collected about Honolulu at low elevations. The coffee, however, was collected on the island of Hawaii, in addition to localities in Honolulu, and much of it came from points from 1,000 to 2,000 feet above sea level. The March collection of coffee came entirely from Kona on the island of Hawaii.

TABLE II.—Percentage of larval parasitism of *Ceratitis capitata* in Hawaii in 1917.—Con.

Host fruit.	Month of collection.	Number of larvae emerging during first 7-6 days.	Percentage of parasitism.				Total.
			<i>Opus humilis</i> .	<i>Diachasma tryoni</i> .	<i>Diachasma tryp.</i>	<i>Tenax gressanus</i> .	
1917.							
Indian almond.....	October.....	6,149	12.1	29.0	.3	15.2	57.2
Do.....	November.....	3,791	12.1	31.5	.5	30.3	74.4
Do.....	December.....	901	11.0	32.3	2.5	19.1	64.9
Mango.....	February.....						
Do.....	June.....	507	2.8	14.0	.7	8.0	26.1
Do.....	July.....	207	3.9	15.0	2.9	12.6	34.4
Do.....	August.....	35				2.9	11.5
Coffee <sup>a</sup> .....	January.....	3,638	59.3	10.6	.8		70.7
Do.....	February.....	501	72.4	6.6	4.4		83.4
Do.....	March.....	1,722	50.0	33.2	5.9		89.1
Do.....	June.....	554	55.4	1.4			56.8
Do.....	August.....	22		0.1	45.5		51.6
Do.....	September.....	3,527	5.9	11.9	33.2	.03	51.03
Do.....	October.....	2,190	3.4	1.0	60.1	.05	70.55
Do.....	November.....	1,541	35.0	5.9	38.4	.5	79.8
Do.....	December.....	176	21.0	3.4	24.4	.6	49.4
Strawberry guava.....	January.....	18	16.7	33.3			50.0
Do.....	March.....	804	20.0	30.0	7.0	.1	57.7
Do.....	April.....	1,172	13.7	39.8	3.2	1.1	57.8
Do.....	May.....	603	12.3	50.7	1.7	1.8	66.5
Do.....	June.....	1,034	10.9	31.9	1.1	4.4	48.2
Do.....	July.....	939	6.1	74.4		.9	81.1
Do.....	August.....	220	11.8	15.5	23.6	12.3	63.2
Do.....	November.....	160		21.3	6.9	30.6	58.8
Do.....	December.....	267		57.7	3.4	4.1	65.2
Black myrobalan.....	January.....	355	8.2	5.6	4.2	34.6	52.6
Do.....	October.....	2,800	.9	.7	.5	3.9	6.0
Peach.....	April.....	203	.8	47.9		2.3	51.0
Do.....	May.....	833	3.6	37.8		9.2	51.5
Do.....	June.....	2,343	8.1	17.0		14.5	39.6
Do.....	July.....	1,391	6.3	18.0		44.8	69.1
Rose-apple.....	May.....	1,021	8.0	13.6		.1	21.7
Do.....	June.....	386	1.6	20.7		1.3	23.6
Do.....	July.....	274	.7	27.7		1.8	30.2
Do.....	August.....	358	.6	29.1		1.1	30.8
Satin-leaf.....	February.....	2,484	18.4	3.4	1.3	.9	24.0
Do.....	March.....	831	37.5	5.4	10.7	1.1	54.7
French cherry.....	January.....	1,258	13.4	19.5	5.9	.8	39.6
Do.....	February.....	290	13.8	31.4	3.8	1.4	50.4
Do.....	March.....	1,320	13.7	23.8	12.3	.5	50.3
Do.....	April.....	351	5.1	24.8	33.0		62.9
Do.....	May.....	346	6.7	9.5	39.0	.6	55.8
Do.....	November.....	143	12.6	6.3	40.6		59.5
Do.....	December.....	1,421	1.1	2.0	4.4	.1	7.6
West Indian medlar.....	June.....	475	5.3	6.7	.2	3.8	16.0
Do.....	July.....	174	.6	1.2		1.7	3.5
Do.....	August.....	478	.2			.2	.4
Do.....	September.....	78	1.3				1.3
Yellow-wood.....	June.....	153		5.2			5.2
Kamani.....	February.....	250		19.6			19.6
Do.....	March.....	94		57.4			57.4
Yellow oleander.....	January.....	7			14.3	57.1	71.4
Do.....	February.....	56	1.8	1.8	8.9	28.6	41.1
Do.....	March.....	17	5.9	5.9		41.2	53.0

<sup>a</sup> The June collection of coffee came from the Waianae Mountains. *Opus humilis* was first established here, but recently *Diachasma tryoni* was liberated.

TABLE II.—Percentage of larval parasitism of *Ceratitis capitata* in Hawaii in 1917—Con.

Host fruit.	Month of collection.	Number of larvae emerging during first 2-6 days.	Percentage of parasitism.				Total.
			<i>Opus humilis.</i>	<i>Dia-chasma tryoni.</i>	<i>Dia-chasma fullawayi.</i>	<i>Tetrastichus giffardianus.</i>	
	1917.						
Yellow oleander.....	July.....	199	.....	13.6	13.6	47.7	74.9
Do.....	August.....	1,679	.4	7.5	14.4	32.0	54.3
Do.....	September.....	294	1.7	2.0	8.5	25.8	38.0
Do.....	October.....	104	.....	.....	25.0	14.4	39.4
Carambola.....	December.....	59	8.5	5.1	.....	6.8	20.4
Chinese orange.....	February.....	68	4.4	5.9	.....	1.5	11.8
Do.....	March.....	188	6.4	.5	1.6	1.1	9.6
Do.....	April.....	210	6.4	14.6	.5	.9	22.4
Do.....	May.....	100	17.0	11.0	.....	.....	28.0
Do.....	June.....	278	13.3	8.3	.....	2.9	24.5
Do.....	December.....	167	1.2	9.6	3.0	4.2	18.0
Ponay.....	June.....	129	.8	1.6	.....	.....	2.4
Do.....	July.....	386	.....	1.3	.2	2.2	3.7
Do.....	August.....	852	.1	26.3	.....	8.0	34.4
Guava.....	January.....	120	1.7	.8	.....	.....	2.5
Do.....	February.....	89	3.4	.....	.....	.....	3.4
Do.....	March.....	100	1.0	.....	.....	3.0	4.0
Do.....	April.....	472	2.1	.2	.4	.2	2.9
Do.....	May.....	91	8.8	.....	.....	.....	8.8
Do.....	July.....	83	6.0	3.6	.....	48.2	57.8
Do.....	August.....	446	1.8	46.9	.2	1.3	50.2
Do.....	November.....	153	.7	.....	.....	.....	.7
Loquat.....	January.....	19	10.5	21.1	31.6	.....	63.2
Do.....	February.....	109	1.8	11.9	62.4	5.5	81.6
Do.....	March.....	441	7.3	6.8	73.5	5.7	93.3
Do.....	December.....	41	4.9	4.9	51.2	22.0	83.0
Kona orange.....	January.....	.....	.....	.....	.....	.....	.....
Do.....	February.....	14	.....	7.1	.....	.....	7.1
Do.....	March.....	39	5.1	.....	2.6	.....	7.7
Do.....	May.....	14	7.1	14.2	7.1	7.1	35.5
Do.....	November.....	64	.....	.....	10.9	51.6	62.5
Mandarin orange.....	November.....	67	3.0	13.4	.....	.....	16.4
Do.....	December.....	50	.....	12.0	.....	4.0	16.0
Sapodilla.....	February.....	10	30.0	.....	.....	.....	30.0
Do.....	March.....	43	16.3	.....	.....	.....	16.3
Do.....	April.....	42	42.9	.....	.....	.....	42.9
White sapote.....	May.....	396	9.1	1.0	.....	1.0	11.1

Thus, as seen in Table III, nearly one-half of all the Mediterranean fruit-fly larvæ developing during the year were destroyed and this is entirely the result of parasitic importations. This achievement, solely due to the efforts of the Territorial Board of Agriculture and Forestry, is worthy of unusual commendation. Insufficient emphasis perhaps has been placed during recent years upon the utility of these parasites. The constant extinction of at least 45 per cent of all worms developing in fruit, apart from the destruction of this pest through other agencies already present in the islands, notable among them being the ant *Pheidole megacephala* Fabricius, without question greatly decreases the infestation of such fruits as the orange (*Citrus sinensis*) and certain

varieties of mango and avocado not ordinarily susceptible to prohibitive infestation except under conditions permitting an unchecked multiplication of the fly. This point has come to light during 1917, and the statement seems justified that a 50 per cent reduction in the numbers of the fly brings little relief to its favored host fruits, but that those fruits classed as unfavored hosts show a marked improvement in the degree of infestation, and some may become almost wholly free from larvæ. The propagation of such fruits and the encouragement of the parasitic method of control would seem to be the most favorable method of contending with this pest in Hawaii.

TABLE III.—Total parasitism, by month, of all larvæ of *Ceratitis capitata* collected in Hawaii during 1917

Month.	Number of larvæ.	Percentage of parasitism.					Total in 1917.	Total in 1916.
		<i>Opius humilis</i> .	<i>Diachasma tryoni</i> .	<i>Diachasma fullawayi</i> .	<i>Tetrastichus giffardianus</i> .			
January.....	6,183	39.0	15.6	2.0	2.4	59.0	6.98	
February.....	4,568	20.0	8.6	3.0	1.3	32.9	19.5	
March.....	5,907	27.1	22.5	12.6	1.3	63.5	14.7	
April.....	2,861	9.0	27.6	5.5	1.2	43.3	37.64	
May.....	4,439	8.8	26.4	3.4	2.3	40.9	26.69	
June.....	5,919	11.7	16.3	.3	7.8	36.1	27.81	
July.....	4,125	3.9	26.6	.9	19.6	51.0	18.52	
August.....	8,726	2.8	16.4	4.7	9.2	33.1	37.5	
September.....	9,047	5.2	31.3	13.2	2.7	52.4	45.2	
October.....	11,309	7.2	16.4	13.2	8.4	45.2	44.3	
November.....	5,919	17.2	22.6	11.6	20.9	72.3	44.3	
December.....	3,142	5.3	16.7	5.2	7.0	34.2	44.1	
Average, 1917.	72,139	12.7	20.3	7.3	7.2	47.5	.....	
Average, 1916.	83,304	17.2	13.3	2.1	.6	.....	33.2	

This 47 per cent reduction in the abundance of the fruit fly in Hawaii serves another purpose well worthy of mention. It is an important help contributing toward reducing the chances of its introduction to the mainland.

As shown elsewhere by the writers, the parasitism by the braconid *Opius humilis* Silvestri has been found highest during the coolest months of the year. Again during 1917 this was observed. The parasitism by *O. humilis* exceeded that of all of the other parasites combined in the months of January and February and was greater than that of any of the others taken separately during March. During the remaining months the parasite *Diachasma tryoni* was more abundant than the *O. humilis*. It is in the cool months of January, February, and March that the other parasites are so retarded that the *O. humilis* is enabled to gain a considerable foothold.



The average parasitism in all fruits by *Opius humilis* in 1915 was well above that of all the other parasites combined; in 1916 the total parasitism by it was 17.2 per cent, and by the others combined was 16 per cent; while during 1917 the other parasites had so reduced the *O. humilis* that the total by it was only 12.7 per cent and by the others combined was 34.8 per cent. This reduction of the *Opius* is proceeding slowly, but the species is not expected to be entirely annihilated. The coffee collections in the Waianae Mountains at the head of the Waianae Valley during 1917 were interesting. This was the only point at which *O. humilis* had been established when the collections were made. In February it was found parasitizing 89.4 per cent of the larvæ in the coffee; and in June 77.5 per cent were parasitized by it.

The parasite *Tetrastichus giffardianus* Silvestri was more abundantly recovered from fruit collections in 1917 than in any preceding year since its liberation in the islands in 1914. It was recovered, however, only in material collected about Honolulu. Its ability to penetrate to the interior of soft fruits broken or containing holes enables it when numerous to parasitize large numbers of larvæ in such fruits as the mango, the orange, or the common guava. The total average parasitism in larvæ from all fruits during the year was materially increased through the work of *T. giffardianus* in such fruits.

The total parasitism by all species during 1917 was 14.3 per cent higher than in 1916. The average infestation of all fruits combined was, however, not strikingly different from that of 1916.

As shown in Table III, the parasitism by *Opius humilis* is 4.5 per cent less than in 1916, while that of *Diachasma tryoni*, *D. fullawayi* Silvestri, and *Tetrastichus giffardianus* was 7.0 per cent, 5.2 per cent, and 6.6 per cent greater, respectively.

## CORN-ROOTROT AND WHEATSCAB

[PRELIMINARY PAPER]

By G. N. HOFFER, A. G. JOHNSON, and D. ATANASOFF

*Cereal Investigations, Bureau of Plant Industry, United States Department of Agriculture*

In the progress of the investigations of rots of the root, stalk, and ear of Indian corn (*Zea mays*) being conducted by one of the authors (Hoffer) certain things have developed which have such an important bearing on certain phases of the wheat scab problem which is being investigated by the other authors that it has seemed desirable to publish a preliminary statement.

Field observations have shown a conspicuously greater abundance of wheat scab in fields where the wheat (*Triticum aestivum*) was grown immediately following corn that was infected with the Fusarium-rot of the root and stalk. This was especially true in Shelby County, Indiana, where wheat, according to a common practice, was sown in standing corn.

A similar condition was also noted in Dane County, Wisconsin, this summer, where spring wheat was grown immediately following a corn crop. Both in Indiana and in Wisconsin under these conditions abundant development of perithecia of *Gibberella* spp. was found on the old cornstalks remaining in these fields. These perithecia were mature and well filled with viable ascospores at the time when the wheat, in all cases observed, was in head.

Water suspensions of these ascospores both from Indiana and from Wisconsin cornstalks gave positive results when used as inocula on wheat heads. The inocula were applied by means of an atomizer spray. In some cases the heads were subsequently covered with glassine bags for three days, and in other cases no coverings were used. In all cases positive infections were obtained, while the controls sprayed with sterile water and similarly treated remained unaffected. The appearance of the infected heads thus artificially inoculated was identical with that of wheat heads naturally infected with scab.

Cultures from *Gibberella* spp. on old cornstalks have also proved virulently parasitic on the roots of corn plants grown both in large, sterile agar tubes and in sterilized pots of soil.

Similar results on both wheat and corn have been obtained by using cultures from naturally infected wheat heads.

The organisms from both sources have also been found to be similar morphologically. In view of the facts developed by this evidence, it seems certain that these are intercrop parasites which are of great impor-

tance in developing control measures for one of the rots of the root, stalk, and ear of corn and for scab of wheat. As both corn and wheat are such highly important food crops, it is imperative that the investigations bearing on any of these disease problems should be pushed forward with utmost vigor at the present time.

While the data are as yet somewhat fragmentary, it seems evident that, in order to lessen the losses from these diseases on corn and wheat, it is necessary to recognize this intercrop parasitism and develop field practices accordingly. In general, the use of the best-adapted, disease-free seed on clean soil should be practiced. The details of control measures for these diseases of corn and wheat are as yet not worked out, and no simple ones are evident. A crop rotation avoiding wheat following diseased corn is undoubtedly important, unless the cornstalks can be cut close to the ground, removed, and the remaining stubble plowed under before the wheat is planted. Badly scabbed wheat should not be used for seed. Ordinary seed treatments will not control wheatscab; hence, only clean seed on carefully prepared clean soil should be used.

ADDITIONAL COPIES  
OF THIS PUBLICATION MAY BE PROCURED FROM  
THE SUPERINTENDENT OF DOCUMENTS  
GOVERNMENT PRINTING OFFICE  
WASHINGTON, D. C.  
AT  
15 CENTS PER COPY  
SUBSCRIPTION PRICE, \$3.00 PER YEAR

▽







# INDEX

	Page		Page
<i>Abelmoschus esculentus</i> host plant of <i>Ascochyta</i>		<i>Apanteles lactecolor</i> —	
<i>abelmoschi</i> .....	207-212	description.....	134-135
<i>Acer saccharinum</i> , susceptibility of seeds to		distribution.....	134-135
desiccation.....	536	economic importance.....	200-201
Acetic acid. See Acid, acetic.		importation.....	192
<i>Achras zapota</i> , food plant of <i>Ceratitix capi-</i>		life history.....	193-199
<i>lata</i> .....	606-608	parasite of <i>Euprostis chrysotheca</i> .....	193-195
Acid—		seasonal history.....	195-199
acetic, occurrence in silage.....	406-408	secondary parasitism.....	201
butyric, occurrence in silage.....	406-408	summer hosts of.....	199-200
lactic, occurrence in silage.....	406-408	<i>Apanteles</i> —	
propionic, occurrence in silage.....	406-408	<i>arge</i> , male and female, taken at light traps.....	110-117
sulphuric, influence on nitrogen-assimi-		<i>vittata</i> , male and female, taken at light	
lating bacteria.....	323-332	traps.....	110-117
Acidity of Silage Made from Various Crops		Aphid. See <i>Pemphigus populi-transversus</i>	
(paper).....	395-409	Arabis—	
Acidity—		potato. See <i>Macrostiphum solanifolii</i> .	
silage.....	395-409	spinach. See <i>Rhopalosiphum persicae</i> .	
soil, influence on growth of legume bacteria	319	spring grain. See <i>Trioptera graminis</i>	
<i>Acrostagnum</i> sp., causal organism of stem		Apple, rose. See <i>Eugenia jambos</i> .	
lesions of <i>Solanum tuberosum</i> .....	213-220	Apple. See <i>Malus sylvestris</i> .	
<i>Aegle Marmelos</i> , resistance to Citrus-canker.....	340	Archips—	
<i>Aeglinae</i> , susceptibility to Citrus-canker.....	340-341	<i>argyrospila</i> , female, at light traps.....	146-148
<i>Aeglopsis Chevalleri</i> , resistance to citrus-		<i>rosaceana</i> , female, at light traps.....	146-148
canker.....	340	Army, A. C., and Garber, R. J. (paper): Varia-	
Aggressin, natural, value in immunization		tion and Correlation in Wheat, with Special	
against blackleg.....	253-262	reference to Weight of Seed Planted.....	339-351
Air, effect on composition of <i>Ipomoea batatas</i>	273-284	<i>Aristoncha albivenosa</i> , female, at light	
Alfalfa. See <i>Medicago sativa</i> .		traps.....	116-149
Alkalinity, soil, influence on growth of legume		Artschwager, E. F. (paper): Anatomy of	
bacteria.....	319	the Potato Plant, with Special Reference	
Almond, Indian. See <i>Terminalia catappa</i> .		to the Ontogeny of the Vascular System.....	221-252
<i>Alternaria</i> spp., causal organisms of stem		<i>Ascochyta abelmoschi</i> —	
lesions of <i>Solanum tuberosum</i> .....	213-220	causal organism of podrot of <i>Abelmoschus</i>	
Alway, F. J., and McDole, G. R. (paper): Vari-		<i>esculentus</i> .....	298-310
ations in the Moisture Content of the Sur-		cultural character.....	311
face Foot of a Loess Soil as Related to the		description.....	299-310
Hygroscopic Coefficient.....	453-480	inoculations with.....	210-211
<i>Amygdalus persica</i> , food plant of <i>Ceratitix</i>		n. sp.....	209-211
<i>capitata</i> .....	605-608	<i>Atalanta citriodes</i> , resistance to Citrus canker.....	341
Anatomy of the Potato Plant, with Special		Atanasoff, D., et al. (paper): Corn-Rootrot	
Reference to the Ontogeny of the Vascular		and Wheat-scab.....	611-612
System (paper).....	221-252	<i>Autographa</i> —	
<i>Andropogon</i> —		<i>biloba</i> , female, at light traps.....	136-140
<i>halepensis sudanensis</i> , syn. <i>Holcus halepen-</i>		<i>simplex</i> , female, at light traps.....	136-140
<i>sis sudanensis</i> .		Availability of Potash in Some Common	
<i>sorghum</i> —		Soil-Forming Minerals—Effect of Lime	
composition.....	176-185	Upon Potash Absorption by Different	
digestibility.....	176-185	Crops (paper).....	297-316
effect of deep tilling in Great Plains on.....	505-517	<i>Avena sativa</i> —	
effect of soil dynamiting in Great Plains		effect of—	
on.....	505-506	deep tilling in Great Plains on.....	515
effect of subsoiling in Great Plains on.....	481-521	subsoiling in Great Plains on.....	484-504
Anthrax—		silage of, acidity.....	400-409
symptomatic, improved methods of immu-		used in experiments with <i>Puccinia graminis</i>	113
nization against.....	253-262	<i>Averrhoa carambola</i> , food plant of <i>Ceratitix</i>	
toxin, symptomatic, concentration.....	263-264	<i>capitata</i> .....	605-608



Azotobacter—	Page	Buckwheat. See <i>Fagopyrum esculentum</i> .	Page
effect of hydrogen-ion concentration on.....	332	Butter fat, percentage in cow's milk.....	67-96
relation of soil reaction to growth.....	265-271	Butyric acid. See Acid, butyric.	
Bacteria—		Cabbage. See <i>Brassica oleracea</i> .	
alfalfa, effect of hydrogen-ion concentration		<i>Caenurgia</i> —	
on.....	329	<i>crassiuscula</i> , female, at light traps.....	136-142
legume—		<i>erecta</i> , female, at light traps.....	136-142
effect of hydrogen-ion concentration on.....	330-331	Calcium—	
formation of nodules by.....	340	carbonate, effect on solubility of potash in	
identification.....	319-323	soil-forming minerals.....	297-316
influence of—		effect on potash absorption by different	
acidity on growth of.....	319	crops.....	297-316
alkalinity on growth of.....	319	nitrate, requirement for buckwheat in cul-	
sodium hydroxide on reproduction of..	323	ture solutions.....	151-175
sulphuric acid on.....	323-328	<i>Calophyllum inophyllum</i> , food plant of	
lupine, effect of concentration of hydrogen-		<i>Ceratitis capitata</i> .....	606-608
ion.....	328-329	Can Biologic Forms of Stemrust on Wheat	
nitrogen-assimilating—		Change Rapidly Enough to Interfere with	
influence of—		Breeding for Rust Resistance? (paper)...	111-124
sodium hydroxide on.....	228-332	Carambola. See <i>Averrhoa carambola</i> .	
sulphuric acid on.....	328-332	<i>Carpocapsa pomonella</i> , female, at light traps	146, 148
nitrogen-fixing, reproduction of.....	323-333	Carruth, F. E., and Withers, W. A. (paper):	
Bark, tan, use as bedding material.....	187-190	Comparative Toxicity of Cottonseed Prod-	
Barley. See <i>Hordeum</i> spp.		ucts.....	425-452
Beet leafhopper. See <i>Eutettix tenella</i> .		Carsner, E., and Stahl, C. F. (paper): Obtain-	
Beet, sugar. See <i>Beta vulgaris</i> .		ing Beet Leafhoppers Nonvirulent as to	
Berg, W. N. (paper): Concentration of Symp-		Curly-Top.....	393-394
tomatic Anthrax (Blackleg) Toxin.....	263-264	<i>Casimiroa edulis</i> —	
<i>Beta vulgaris</i> , curly-top infection through		food plant of <i>Ceratitis capitata</i> .....	606-608
<i>Eutettix tenella</i> .....	393-394	resistance to Citrus-canker.....	340
Bethel, E., et al. (paper): Piñon Blister		Cattle—	
Rust.....	411-424	occurrence of coccidioid granuloma in.....	533-542
Beyer, A. H., and Luginbill, P. (paper):		value of Sudan-grass hay as feed for.....	176-185
Contribution to the Knowledge of Toxop-		Cauliflower. See <i>Brassica oleracea botrytis</i> .	
tere graminum in the South.....	97-110	<i>Ceratitis capitata</i> , parasitism in Hawaii in	
Biotite, availability of potash in.....	297-316	1917.....	605-610
Black myrobalan. See <i>Terminalia chebula</i> .		<i>Chaetochytrium glutinosa</i> , resistance to Citrus-	
Blackleg—		canker.....	341
improved methods of immunization		<i>Chalcas exotica</i> , resistance to Citrus-canker...	340
against.....	253-264	Cherry—	
toxin, concentration of.....	263-264	French. See <i>Eugenia uniflora</i> .	
Blister-rust, piñon, caused by <i>Cronartium</i>		wild. See <i>Prunus</i> spp.	
<i>occidentale</i> .....	411-424	Chilcott, E. C., and Cole, J. S. (paper): Sub-	
Blood, rat, recovery of trypanosomes from.....	573-576	soiling, Deep Tilling, and Soil Dynamiting	
Bluegrass, Kentucky. See <i>Poa pratensis</i> .		in the Great Plains.....	481-521
<i>Botrytis</i> sp., casual organism of stem lesions of		Chinese orange. See <i>Citrus japonica</i> .	
<i>Solanum tuberosum</i> .....	213-220	<i>Chloridea obsoleta</i> , female, at light traps....	147, 148
Brassica—		<i>Chrysophyllum olivaeforme</i> , food plant of	
<i>napa</i> , food-plant of <i>Pemphigus populi-</i>		<i>Ceratitis capitata</i> .....	606-608
<i>transversus</i> .....	585	Cictrange, resistance to Citrus-canker.....	350
<i>nigra</i> , food plant of <i>Pemphigus populi-</i>		<i>Cirphis unipuncta</i> , female, at light traps....	136-143
<i>transversus</i> .....	585	Citridia, resistance to Citrus-canker.....	349
<i>oleracea</i> —		Citrindarin—	
<i>botrytis</i> , food plant of <i>Pemphigus populi-</i>		overwintering of Citrus-canker organism	
<i>transversus</i> .....	585	in.....	523-524
<i>capitata</i> , food plant of <i>Pemphigus populi-</i>		resistance to Citrus-canker.....	350
<i>transversus</i> .....	577-594	Citrange—	
<i>gemmifera</i> , food-plant of <i>Pemphigus</i>		resistance to Citrus-canker.....	348-349
<i>populi-transversus</i> .....	585	Rusk, overwintering of Citrus-canker organ-	
host of <i>Plasmodiophora brassicae</i> .....	543-572	ism in.....	523-524
tissue invasion of by <i>Plasmodiophora</i>		Savage, overwintering of Citrus-canker organ-	
<i>brassicae</i> .....	543-572	anism in.....	523-524
<i>rapa</i> , food plant of <i>Pemphigus populi-</i>		Citranequat, resistance to Citrus-canker....	350
<i>transversus</i> .....	578	Citranguma, resistance to Citrus-canker.....	350
Broccoli. See <i>Brassica oleracea botrytis</i> .		Citrease, susceptibility to Citrus-canker.....	340-341
Broom corn. See <i>Andropogon sorghum</i> .		Citriacae, susceptibility to Citrus-canker.....	341
Brussels sprouts. See <i>Brassica oleracea</i>		<i>Citropis Schweinfurthii</i> , resistance to Citrus-	
<i>gemmifera</i> .		canker.....	341

	Page		Page
Citrumelo, resistance to Citrus-canker.....	349	Corn—	
<i>Citrus</i> —		brum. See <i>Andropogon scoparius</i> .	
<i>aurantifolia</i> , resistance to Citrus-canker.....	346	See also <i>Zea mays</i> .	
<i>exaltis</i> , resistance to Citrus-canker.....	348	Corn-Rootrot and Wheatsoy (paper).....	611-612
genus, susceptibility of wild relatives of, to		<i>Cornus didymus</i> , food plant of <i>Pemphigus</i>	
Citrus-canker.....	346-355	<i>populi-transversus</i> .....	583
<i>grandis</i> —		Correlation between the Percentage of Fat in	
overwintering of Citrus-canker organism		Cow's Milk and the Yield (paper).....	67-90
in.....	523-524	Cotton. See <i>Gossypium</i> spp.	
resistance to Citrus-canker.....	346	Cottonseed—	
hybrid. See Hybrid, citrus.		effect of cooking on toxicity.....	426-429
<i>Hystrix</i> , resistance to Citrus-canker.....	344	meal, toxicity.....	425-431
<i>japonica</i> , food plant of <i>Ceratitidis capitata</i> .....	606-608	toxicity.....	425-431
<i>Medica</i> , resistance to Citrus-canker.....	344	Cottonwood. See <i>Populus</i> spp.	
<i>mitis</i> , resistance to Citrus-canker.....	347	Cow—	
<i>nobilis</i> —		milk of, correlation between percentage of	
<i>deliciosa</i> , food plant of <i>Ceratitidis capitata</i> .....	606-608	fat and yield.....	67-90
resistance to Citrus-canker.....	347	value of Sudan-grass hay as feed for.....	120-185
var. <i>unihis</i> , overwintering of Citrus-		Cowpea. See <i>Vigna sinensis</i> .	
canker organism in.....	523-524	<i>Crambus hortuella</i> , female, at light traps.....	142-148
<i>sinensis</i> , food plant of <i>Ceratitidis capitata</i> .....	606-608	Crocker, W., and Harrington, G. T. (paper)	
spp. resistance to Citrus-canker.....	344-348	Resistance of Seeds to Desiccation.....	613-614
susceptibility of wild relatives of, to Citrus-		<i>Cronarium</i> —	
canker.....	346-344	<i>occidentale</i> —	
Citrus-canker—		cause of pith blister-rust.....	441-444
organism of, overwintering.....	523-524	comparison with <i>Cronarium rhizoda</i> .....	414-415
resistance of Citrus plants to.....	337-357	description.....	411-415
susceptibility of Citrus plants to.....	337-357	dissemination.....	406-409
<i>Claucaena</i> —		distribution.....	415-418
<i>lansium</i> , resistance to Citrus-canker.....	340	effect on hosts.....	412-414
<i>wampi</i> , food plant of <i>Ceratitidis capitata</i> .....	606-608	n. sp.....	413-415
<i>Clonostachys</i> sp., casual organism of stem le-		overwintering.....	430-432
sions of <i>Solanum tuberosum</i> .....	213-220	inoculation.....	418-422
Clubroot, disease of crucifers caused by <i>Plas-</i>		<i>Ribicola</i> , comparison with <i>Cronarium occi-</i>	
<i>modiophora brassicae</i> .....	543-572	<i>dentale</i> .....	414-415
Coccidioid granuloma, occurrence in—		Cruciferae, occurrence of galls in.....	572-594
cattle.....	533-542	Damping off, resemblance to whitespot dis-	
man.....	533-534	ease of plants.....	595
<i>Coccidioides immitis</i> —		<i>Datana ministra</i> , female, at light traps.....	136-144
casual organism of coccidioid granuloma.....	533-542	Davenport, A., and Fred, E. B. (paper): In-	
cultural characters.....	535-536	fluence of Reaction on Nitrogen-Assimilat-	
description.....	534-536	ing Bacteria.....	317-336
inoculations with.....	536-540	Deep tilling of soil, effect in Great Plains.....	506-517
life cycle.....	534-535	Desiccation, resistance of seeds to.....	515-532
<i>Coffea arabica</i> , food plant of <i>Ceratitidis capitata</i>	606-608	<i>Diachasma</i> —	
Coffee. See <i>Coffea arabica</i> .		<i>fullawayi</i> , parasite of <i>Ceratitidis capitata</i> .....	606-610
Cole, J. S., and Chilcott, E. C. (paper): Sub-		<i>tryoni</i> , parasite of <i>Ceratitidis capitata</i> .....	606-610
soiling, Deep Tilling, and Soil Dynamiting		<i>Diaeris virginica</i> , female, at light traps.....	136-138
in the Great Plains.....	481-521	Dog, susceptibility to coccidioid granul-	
Comparative Study of Salt Requirements for		oma.....	516-538
Young and for Mature Buckwheat Plants in		Drouth, relation to moisture content of loess	
Solution Cultures, A (paper).....	151-175	soil.....	457-461
Comparative Toxicity of Cottonseed Products		Dynamiting of soil, effect in Great Plains.....	505-517
(paper).....	425-452	Edson, H. A., and Shapovalov, M. (paper):	
Complement fixation, improved method for		Potato-Stem Lesions.....	273-220
recovering trypanosomes from blood of rats		Effect of Different Oxygen Pressures on the	
for antigen purposes.....	573-576	Carbohydrate Metabolism of the Sweet	
Composition and Digestibility of Sudan-Grass		Potato (paper).....	273-284
Hay (paper).....	176-185	<i>Elasmopalpus lignosellus</i> , female, at light	
Concentration of Symptomatic Anthrax		traps.....	145-148
(Blackleg) Toxin (paper).....	263-264	Emmer. See <i>Triticum dicoccum</i> .	
Conifers, whitespot disease of.....	595-604	<i>Eremocirus glauca</i> , resistance to Citrus-	
Contribution to the knowledge of Toxoptera		canker.....	342
graminum in the South (paper).....	97-110	<i>Eriobotrya japonica</i> , food plant of <i>Ceratitidis</i>	
<i>Corethrobia</i> sp., casual organism of stem le-		<i>capitata</i> .....	606-608
sions of <i>Solanum tuberosum</i> .....	216-219	Errata.....	IV

	Page		Page
<i>Erysiphe graminis</i> DC., biologic forms of, on genus <i>Bromus</i> broken down by use of bridging hosts.....	112	Garber, R. J., and Army, A. C. (paper): Variation and Correlation in Wheat, with Special Reference to Weight of Seed Planted.....	359-392
<i>Estigmene acrea</i> , female, at light traps.....	136-138	"Germ-free vaccine." See Vaccine, germ-free.	
<i>Eugenia</i> —		<i>Gibberella</i> spp. on old cornstalks.....	611
<i>jambos</i> , food plant of <i>Ceratitis capitata</i> .....	606-608	Gillette, C. P. (paper): Descriptions of Stages of <i>Pemphigus populi-transversus</i> and Its Gall.....	589-592
<i>uniflora</i> , food plant of <i>Ceratitis capitata</i> .....	606-608	Giltner, L. T. (paper): Occurrence of Coccidioid Granuloma (Oidiomycosis) in Cattle.....	533-542
<i>Euproctis chrysorrhoea</i> —		<i>Glycosmis pentaphylla</i> , resistance to Citrus-canker.....	340
distribution.....	191	<i>Gossypium</i> spp., effect of subsoiling in Great Plains on.....	484-504
female, at light traps.....	145, 148	Gossypol, toxicity.....	425-452
host of—		<i>Gramineae</i> spp., resistance of seeds to desiccation.....	526-532
<i>Apanteles lacteicolor</i> .....	191-206	Grapefruit. See <i>Citrus grandis</i> .	
<i>Meteorus versicolor</i> .....	191-206	Grass—	
life history.....	191-192	blue, Kentucky. See <i>Poa pratensis</i> .	
parasite of—		Johnson. See <i>Holcus halepensis</i> .	
<i>Malus sylvestris</i> .....	191-192	Sudan—	
<i>Prunus</i> spp.....	191-192	composition of hay.....	176-185
<i>Pyrus communis</i> .....	191-192	digestibility of hay.....	176-185
<i>Quercus</i> spp.....	191-192	yield for hay.....	176
<i>Eutetrix tenella</i> —		See <i>Holcus halepensis sudanensis</i> .	
nonvirulent as to curly-top.....	393-394	Great Plains—	
transmission of curly-top of <i>Beta vulgaris</i> by.....	393-394	effect of—	
<i>Fagopyrum esculentum</i> , salt requirements in culture solutions.....	151-175	deep tilling in.....	505-517
Faustrime, resistance to Citrus-canker.....	348	soil dynamiting in.....	505-517
Faustrimon, resistance to Citrus-canker.....	348	subsoiling in.....	481-521
Feed—		<i>Grossularia</i> spp.—	
cow's, value of Sudan-grass hay as.....	176-185	injury by <i>Cronartium occidentale</i> .....	422-423
poultry, Southern, mineral content of.....	125-134	inoculation with <i>Cronartium occidentale</i> .....	419-420
<i>Feltia</i> spp., female, at light traps.....	136-144	Guava—	
Female Lepidoptera at Light Traps (paper).....	135-149	strawberry. See <i>Psidium cattleianum</i> .	
<i>Feronia Limonia</i> , resistance to Citrus-canker.....	341	See also <i>Psidium guajava</i> .	
<i>Feroniella lucida</i> , resistance to Citrus-canker.....	341	Guinea pig, susceptibility to coccidioid granuloma.....	536-537
Feroniinae, susceptibility to Citrus-canker.....	341	Gypsum, effect on water-soluble potash.....	61-66
Filtrate, toxic culture, value in immunization against blackleg.....	261-262	<i>Halimolobos</i> —	
Flax. See <i>Linum usitatissimum</i> .		<i>caryae</i> , female, at light traps.....	147-148
Fly, fruit, Mediterranean. See <i>Ceratitis capitata</i> .		<i>tesellaris</i> , female, at light traps.....	136-139
<i>Fortunella</i> —		Hardy, J. I. (paper): Influence of Humidity upon the Strength and the Elasticity of Wool Fiber.....	285-296
<i>crassifolia</i> , resistance to Citrus-canker.....	342-343	Harrington, G. T., and Crocker, W. (paper): Resistance of Seeds to Desiccation.....	515-532
<i>hindii</i> , resistance to Citrus-canker.....	343	Harter, L. L. (paper): A Hitherto Unreported Disease of Okra.....	207-212
<i>margarita</i> , resistance to Citrus-canker.....	342	Hartley, C. (paper): Stem Lesions Caused by Excessive Heat.....	595-604
Fowl—		Hasselbring, H. (paper): Effect of Different Oxygen Pressures on the Carbohydrate Metabolism of the Sweet Potato.....	273-284
mineral requirements of.....	125-134	Hawaii, Mediterranean fruit fly during 1917 in.....	605-610
See Poultry.		Hedgecock, G. C., Bethel, E., and Hunt, N. R. (paper): Piñon Blister-Rust.....	411-424
Fred, E. B., and Davenport, A. (paper): Influence of Reaction on Nitrogen-Assimilating Bacteria.....	317-336	<i>Helianthus annuus</i> , effect of deep tilling in Great Plains on.....	508-517
French cherry. See <i>Eugenia uniflora</i> .		<i>Hemileuca oliviae</i> , female, at light traps.....	146, 148
Fruit, Citrus, susceptibility to Citrus-canker.....	344-348		
Fruit fly, Mediterranean. See <i>Ceratitis capitata</i> .			
<i>Fusarium</i> spp., causal organisms of stem lesions of <i>Solanum tuberosum</i> .....	213-220		
<i>Fusarium-rot</i> on corn.....	611		
Gaessler, W. G., and McCandlish, A. C. (paper): Composition and Digestibility of Sudan-Grass Hay.....	176-185		
Gainey, P. L. (paper): Soil Reaction and the Growth of <i>Azotobacter</i> .....	265-272		
Galls occurrence in—			
crucifers.....	577-594		
<i>Populus</i> spp.....	577-594		

	Page		Page
<b>Hen—</b>		Katipp, B. F. (paper): Mineral Content of Southern Poultry Feeds and Mineral Requirements of Growing Fowls	325-334
effect of cottonseed products on—		Kelser, R. A. (paper): Improved Methods of Immunization Against Symptomatic Anthrax (Blackleg)	253-262
egg production of	441-442	Kentucky bluegrass. See <i>Poa pratensis</i> .	
growth of	438-442	Kona orange. See <i>Citrus sinensis</i> .	
<b>Hesperethusa crenulata</b> , resistance to Citrus-canker	341	Kunkel, L. O. (paper): Tissue Invasion by <i>Plasmidiophora brassicae</i>	541-552
Hoffer, G. N., Johnson, A. G., and Atanasoff, D. (paper): Corn-Rootrot and Wheat-scab	611-612	Lactic acid, occurrence in silage	408-408
<b>Holcus—</b>		Lavanginae, susceptibility to Citrus canker	341
<i>halepensis</i> resistance of seeds to desiccation	527-532	Leaf, satin. See <i>Chrysophyllum discolorum</i> .	
<i>sudanensis</i> , resistance of seeds to desiccation	527-532	Leafhopper, beet. See <i>Eutettix tenella</i> .	
Honolulu, parasitism of <i>Tetrastichus giffard-</i>	606-610	Leaves, oak, use as bedding material	187-190
<i>ianus</i> , near	606-610	Legume, bacteria. See Bacteria, legume.	
<b>Hordeum—</b>		Legumes, influence of reaction of soil on growth of	317-320
<i>jubatum</i> , used in experiments with <i>Puccinia graminis tritici-compacti</i>	117	<i>Lepidium virginicum</i> , food plant of <i>Pemphigus populi-transversus</i>	585
spp., effect of subsoiling in Great Plains on	484-504	<i>populi-transversus</i>	585
<i>compacti</i>	117-121	Lepidoptera, female, at light traps	115-149
host plant of <i>Puccinia graminis tritici-compacti</i>	117-121	Lesions, stem, of <i>Solanum tuberosum</i> —	
host plant of <i>Puccinia graminis tritici-compacti</i>	117-121	casual organisms of	211-219
resistance of seeds to desiccation	527-532	inoculations with fungi from	214-219
<b>Hitherto Unreported Disease of Okra</b> , (paper)	207-212	isolations of fungi from	211-214
Humidity, influence on wool fiber	285-295	Life History of <i>Pemphigus populi-transversus</i> (paper)	577-594
Hybrid, Citrus, overwintering of Citrus-canker organism in	523-524	<b>Ligniera—</b>	
Hybrids, Citrus, susceptibility to Citrus-canker	348-354	<i>graminis</i> , tissue invasion by	560
<b>Hydrogen-ion—</b>		<i>junci</i> , tissue invasion by	560
concentration of, effect on—		Lime. See Calcium.	
<i>allia</i> bacteria	329	Limelo, resistance to Citrus-canker	350
Azotobacter	332	Limequat, resistance to Citrus-canker	350
lupine bacteria	328-329	<i>Linum catharticum</i> , effect of subsoiling in Great Plains on	484-504
Hygrosopic coefficient, relation of moisture content of loess soil to	453-480	Loquat. See <i>Eriobotrya japonica</i> .	
Improved Method for Recovering Trypanosomes from the Blood of Rats for Antigen Purposes in Connection with Complement Fixation, An (paper)	573-576	Luginbill, P., and Beyer, A. H. (paper): Contribution to the Knowledge of Toxoptera graminum in the South	97-110
Improved Methods of Immunization against Symptomatic Anthrax (Blackleg) (paper)	253-262	McCandlish, A. C., and Gaessler, W. G. (paper): Composition and Digestibility of Sudan-Grass Hay	176-185
<b>Indian—</b>		McClintock, J. A., and Smith, L. B. (paper): True Nature of Spinach-Blight and Relation of Insects to Its Transmission	1-60
almond. See <i>Terminalia catappa</i> .		McDole, G. R., and Alway, F. J. (paper): Variations in the Moisture Content of the Surface Foot of the Loess Soil as Related to the Hygrosopic Coefficient	453-480
corn. See <i>Zea mays</i> .		McMiller, P. R. (paper): Influence of Gypsum upon the Solubility of Potash in Soils	61-66
Influence of Gypsum upon the Solubility of Potash in Soils (paper)	61-66	<i>Macrosporum udanifolii</i> , agency in dissemination of spinach-blight	11-56
Influence of Humidity upon the Strength and the Elasticity of Wool Fiber (paper)	285-296	Magnesium sulphate, requirement for buckwheat in culture solutions	151-175
Influence of Reaction on Nitrogen-Assimilating Bacteria (paper)	317-336	<i>Malus sylvestris</i> , food plant of <i>Euprodia chrysorrhoea</i>	192-192
<b>Ipomoea batatas—</b>		Man, occurrence of coccidoidal granuloma in	533-534
effect of—		Mandarin orange. See <i>Citrus nobilis deliciosa</i> .	
air on composition of	273-284	<i>Mangifera indica</i> , food plant of <i>Ceratit capitata</i>	606-608
hydrogen on composition of	273-284	Mango. See <i>Mangifera indica</i> .	
oxygen on composition of	273-274	Manure, conservation, in relation to bedding materials	187-190
<i>Isia isabella</i> , female, at light traps	136-139		
Johnson, A. G., et al. (paper): Corn-Rootrot and Wheat-scab	611-612		
Johnson grass. See <i>Holcus halepensis</i> .			
Jones, T. H. (paper): Life History of <i>Pemphigus populi-transversus</i>	577-594		
Kafir. See <i>Andropogon sorghum</i> .			
Kamani. See <i>Calophyllum inophyllum</i> .			

	Page		Page
Maple, silver. See <i>Acer saccharinum</i> .		Obtaining Beet Leafhoppers Nonvirulent as to Curly-Top (paper).....	393-394
Martin, W. H., and Shive, J. W. (paper): A Comparative Study of Salt Requirements for Young and for Mature Buckwheat Plants in Solution Cultures.....	151-173	Occurrence of Coccidioid Granuloma (Oidomycosis) in Cattle (paper).....	513-542
<i>Medicago sativa</i> , silage of, acidity.....	395-409	<i>Ocrotia elliptica</i> , food plant of <i>Ceratitis capitata</i> .....	606-608
Mediterranean fruit fly. See <i>Ceratitis capitata</i> .		Oidionmycosis occurrence in—	
Medlar, West Indian. See <i>Mimusops elengi</i> .		cattle.....	533-542
<i>Melana diffusa</i> , female, at light traps.....	136-140	man.....	533-534
<i>Meteorus versicolor</i> —		Okra. See <i>Abelmoschus esculentus</i> .	
description.....	201-205	Oleander, yellow. See <i>Thevetia nerifolia</i> .	
distribution.....	192-194	<i>Opus humilis</i> , parasite of <i>Ceratitis capitata</i> .	606-610
economic importance.....	205	Orange—	
importation.....	192	Chinese. See <i>Citrus japonica</i> .	
life history.....	202-204	kona. See <i>Citrus sinensis</i> .	
parasite of <i>Euprocitis chrysorrhoea</i> .....	191-206	mandarin. See <i>Citrus nobilis deliciosa</i> .	
seasonal history.....	202-204	Satsuma. See <i>Citrus nobilis</i> var. <i>unshiu</i> .	
summer hosts of.....	204-205	Orthoclase, availability of potash in.....	297-316
<i>Microcitrus</i> —		<i>Ostenfeldella diplanthera</i> , tissue invasion by.....	569
australasia, resistance to Citrus-canker.....	343	Overwintering of the Citrus-Canker Organism in the Bark Tissue of Hardy Citrus Hybrids (paper).....	523-524
australis, resistance to Citrus-canker.....	343-344	Oxygen, effect on composition of <i>Ipomoea batatas</i> .....	273-284
<i>Garrowayi</i> , resistance to Citrus-canker.....	343	<i>Panicum miliaceum</i> , effect of deep tilling in Great Plains on.....	514-516
<i>Microcline</i> , availability of potash in.....	297-316	Parasitism, intercrop, control of.....	610
Milk, cow's—		Parker, John H., et al. (paper): Can Biologic Forms of Stemrust on Wheat Change Rapidly Enough to Interfere with Breeding for Rust Resistance?.....	111-123
correlation between percentage of fat and yield.....	67-96	Peach. See <i>Amygdalus persica</i> .	
value of—		Pear. See <i>Pyrus communis</i> .	
alfalfa hay in production of.....	178-179	Peas. See <i>Pisum</i> spp.	
Sudan-grass hay in production of.....	178-179	Peat, use as bedding material.....	187-190
Millet. See <i>Panicum miliaceum</i> .		Peltier, G. L., and Neal, D. C. (paper): Overwintering of the Citrus-Canker Organism in the Bark Tissue of Hardy Citrus Hybrids.....	523-524
Milo. See <i>Andropogon sorghum</i> .		Peltier, G. L. (paper): Susceptibility and Resistance to Citrus-Canker of the Wild Relatives Citrus Fruits, and Hybrids of the Genus Citrus.....	317-358
<i>Mimusops elengi</i> , food plant of <i>Ceratitis capitata</i> .....	606-608	Pemberton, C. E., and Willard, H. F. (paper): Work and Parasitism of the Mediterranean Fruit Fly in Hawaii during 1917.....	605-610
Mineral Content of Southern Poultry Feeds and Mineral Requirements of Growing Poultry (paper).....	125-134	<i>Pemphigus</i> —	
Moisture, variations in loess soil.....	453-480	<i>populicautis</i> , galls of.....	592
Mold, vegetable, use as bedding material.....	187-190	<i>populi-rumulorum</i> , galls of, comparison with <i>Pemphigus populi-transversus</i> .....	592
Moth—		<i>populi-transversus</i> —	
brown-tail—		description of gall of.....	584, 589-592
parasites of.....	191-206	description of stages.....	589, 592
See <i>Euprocitis chrysorrhoea</i> .		distribution.....	580-581
See also Lepidoptera.		formation of gall by.....	587
Muesebeck, C. F. W. (paper): Two Important Introduced Parasites of the Brown-Tail Moth.....	191-206	host plants of.....	580-581
Muscovite, availability of potash in.....	297-316	injury caused by.....	584-586
Mustard. See <i>Brassica nigra</i> .		life history.....	577-594
Myrobalan, black. See <i>Terminalia chebula</i> .		migration.....	581-586
"Natural aggression." See Aggression, natural.		occurrence of winged migrants in soil.....	586
Neal, D. C., and Peltier, G. L. (paper): Overwintering of the Citrus-Canker Organism in the Bark Tissue of Hardy Citrus Hybrids.....	523-524	oviposition.....	587
Neidig, R. E. (paper): Acidity of Silage Made from Various Crops.....	395-409	reproduction.....	586-587
Nitrogen—		seasonal history.....	587-589
effect on composition of <i>Ipomoea batatas</i> .....	273-284	<i>Peridermium</i> —	
fixation by bacteria, influence of reaction on.....	317-336	occidentale, aecial stage of <i>Cromartium occidentale</i> .....	413-418
Nitrogen-assimilating bacteria, influence of reaction on.....	317-336	<i>strobi</i> , aecial stage of <i>Cromartium occidentale</i> .....	413-415
Noctua c-nigrum, female, at light traps.....	136-144		
Nodules, formation from legume bacteria.....	320		
Oak. See <i>Quercus</i> spp.			
Oats. See <i>Avena sativa</i> .			

	Page		Page
<i>Pheidole megacephala</i> Fabricius, native parasite of fruit pests in Hawaii.....	608	<i>Populus</i> —Continued	
<i>Phoma</i> sp., causal organism of stem lesions of <i>Solanum tuberosum</i> .....	216-219	<i>deltoidea</i> —	
<i>Phthorimaea operculella</i> , female, at light traps.....	145-148	food plant of <i>Pemphigus popule-transversus</i> .....	580-581
Piemeisel, F. J., et al. (paper): Can Biologic Forms of Stemrust on Wheat Change Rapidly Enough to Interfere with Breeding for Rust Resistance?.....	111-113	formation of galls on.....	581
Pig, toxicity of cottonseed products to.....	447-449	percentage of leaf petioles showing galls.....	581
Pine. See <i>Pinus</i> spp.		<i>monilifera</i> —	
Piñon Blister-Rust (paper).....	421-424	food plant of <i>Pemphigus popule-transversus</i> .....	580-581
Pinon. See <i>Pinus edulis</i> ; <i>Pinus monophylla</i> .		syn. <i>Populus deltoidea</i> .....	580-581
<i>Pinus</i> —		spp., occurrence of galls in.....	577-584
<i>edulis</i> , host plant of <i>Cronartium occidentale</i> .....	411-424	<i>Portheia dispar</i> , female, at light traps.....	147-148
<i>monophylla</i> , host plant of <i>Cronartium occidentale</i> .....	411-424	Potash—	
spp.—		solubility in soils, effect of gypsum on.....	61-66
resemblance of damping-off to whiteness.....	595	See Potassium.	
stem lesions caused by bending.....	601-602	Potassium—	
stem lesions caused by heat.....	595	availability in soil-forming minerals.....	297-316
whiteness disease of.....	595-601	phosphate, requirement for buckwheat in culture solutions.....	330-335
<i>Pisum</i> spp., silage of, acidity.....	400-409	sulphate, availability of potash in.....	297-316
Plains, Great. See Great Plains.		Potato—	
Plant, rutaceous, susceptibility to Citrus-canker.....	342-344	sweet. See <i>Ipomoea batatas</i> .	
<i>Plasmodiophora brassicae</i> —		See also <i>Solanum tuberosum</i> .	
galls, comparison with <i>Spongospora subterranea</i> .....	567-568	Potato-Stem Lesions (paper).....	411-420
histology of club caused by.....	557-561	Poultry, effect of cottonseed products on—	
infection of young roots by.....	561-562	egg production.....	441-442
morphology of club caused by.....	546-547	growth.....	448-449
nature of infection by.....	545-546	Propionic acid, occurrence in silage.....	466-468
parasite of <i>Brassica oleracea</i> .....	543-572	<i>Prunus</i> spp., food plant of <i>Euproctis chrysorrhoea</i> .....	191-192
passage through host tissues.....	554-556	<i>rhoea</i> .....	191-192
production of branch roots and shoots caused by.....	556-557	<i>Pseudomonas citri</i> , inoculation of Citrus plants with.....	637-637
relation to host tissues.....	562-567	<i>Psidium</i> —	
stages in club formation caused by.....	547-554	<i>cattianum</i> , food plant of <i>Ceratitus capitata</i> .....	660-668
time of infection by.....	545-546	<i>quajaro</i> , food plant of <i>Ceratitus capitata</i> .....	660-668
tissue invasion by.....	543-572	<i>Puccinia</i> —	
<i>Plasmodiophoraceae</i> , tissue invasion by.....	543-572	<i>dispersa</i> Erikss., reference to Ward's study of.....	112
Plummer, J. K. (paper): Availability of Potash in Some Common Soil-Forming Minerals—Effect of Lime Upon Potash Absorption by Different Crops.....	297-316	<i>glumarum</i> Erikss. and Henn., resistance in wheats to a recessive Mendelian character.....	112
<i>Phutella maculipennis</i> , female, at light traps.....	146, 148	<i>graminis</i> Pers., whether bridging hosts can be used with, to infect plants normally immune.....	112
<i>Poa pratensis</i> , resistance of seeds of to desiccation.....	526-532	<i>tritici</i> —	
Podspot of <i>Adelmoschus esculentus</i> —		effect of wheat hybrids on.....	117-121
cause.....	208-210	inoculations with.....	117-121
description.....	208	<i>Pyrus communis</i> , food plant of <i>Euproctis chrysorrhoea</i> .....	191-192
origin.....	207-208	<i>Quercus</i> spp., food plant of <i>Euproctis chrysorrhoea</i> .....	191-192
Poisoning, cottonseed.....	425-452	Rabbit—	
<i>Polia renigera</i> , female, at light traps.....	136-141	susceptibility to coccidial granuloma.....	536-538
<i>Poncirus trifoliata</i> —		toxicity of cottonseed products to.....	415-418
overwintering of Citrus-canker organism in.....	523-524	Rape. See <i>Brassica napus</i> .	
resistance to Citrus-canker.....	522	Rat—	
Poplar. See <i>Populus</i> spp.		recovery of trypanosomes from blood of.....	573-576
<i>Populus</i> —		toxicity of cottonseed products to.....	427-435
<i>balsamifera</i> , food plant of <i>Pemphigus popule-transversus</i> .....	580-582	Resistance of Seeds to Desiccation (paper).....	525-527
		Reynolds, F. H., and Schoening, H. W. (paper): An Improved Method for Recovering Trypanosomes from the Blood of Rats for Antigen Purposes in Connection with Complement Fixation.....	573-576
		<i>Rhizoctonia</i> spp., casual organisms of stem lesions of <i>Solanum tuberosum</i> .....	213, 220

	Page		Page
<i>Rhizodium leguminosarum</i> , effect on reaction of culture medium.....	332	Soil—Continued	
<i>Rhopalosiphum persicae</i> , agency in dissemination of spinach blight.....	11-56	reaction of—	
<i>Ribes</i> spp.—		influence on legumes.....	
injury by <i>Cronartium occidentale</i> .....	422-423	relation to growth of <i>Azotobacter</i> .....	265-271
inoculation with <i>Cronartium occidentale</i> .....	419-420	subsoiling, effect in Great Plains.....	481-521
Rice, wild. See <i>Zizania palustris</i> .		Soil Reaction and the Growth of <i>Azotobacter</i> (paper).....	265-271
Roberts, E. (paper): Correlation between the Percentage of Fat in Cow's Milk and the Yield.....	67-96	<i>Solanum tuberosum</i> —	
<i>Roripa</i> sp., food plant of <i>Pemphigus populi-transversus</i> .....	583	anatomy.....	221-236
Rose-apple. See <i>Eugenia jambos</i> .		gross morphology.....	223-225
<i>Ruphia rostellata</i> , host of <i>Tetramyxa parasitica</i> .....	569	ontogeny.....	237-250
Rust—		secondary growth.....	249-250
blister, pñon. See Blister-rust, pñon.		stem lesions of.....	213-220
stem. See <i>Puccinia graminis tritici</i> . See also <i>Puccinia graminis tritici-compacti</i> .		<i>Sorghum halepense</i> , syn. <i>Holcus halepensis</i> .	
Rutaceous plants. See Plant, rutaceous.		Sorghum. See <i>Andropogon sorghum</i> .	
Rye. See <i>Secale cereale</i> .		<i>Sorodiscus callitrichis</i> , tissue invasion by.....	569
<i>Salix</i> spp., susceptibility to desiccation.....	526	<i>Sororphaera veronica</i> , tissue invasion by.....	568
<i>Sanninoidea opalecens</i> , female, at light traps 146-148		South, <i>Toxoptera graminum</i> in.....	97-110
Sapodilla. See <i>Ackras sapota</i> .		Soybean meal, toxicity.....	428-448
Sapote, white. See <i>Casimiroa edulis</i> .		Species, new.....	209-211, 413-415
Satin-leaf. See <i>Chrysophyllum pluvialeforme</i> .		<i>Spinacea oleracea</i> , blight of—	
Satsuma orange. See <i>Citrus nobilis</i> var. <i>unshiu</i> .		cause of dissemination.....	6
Sawdust, use as bedding material.....	187-190	control.....	55-56
Schoening, H. W., and Reynolds, F. H. (paper): An Improved Method for Recovering Trypanosomes from the Blood of Rats for Antigen Purposes in Connection with Complement Fixation.....	573-576	description.....	2-4
<i>Sclerotinia</i> sp., casual organism of stem lesions of <i>Solanum tuberosum</i> .....	213-220	history.....	4-5
<i>Secale cereale</i> , whiteness disease of.....	598	insect transmission.....	10-53
Seel, resistance to desiccation.....	525-532	relation of bacteria to.....	7-8
<i>Severinia busifolia</i> , resistance to Citrus-canker 341		relation of drainage to.....	6-7
Shapovalov, M., and Edson H. A. (paper): Potato-Stem Lesions.....	213-220	relation of fertilizer to.....	7
Shavings, use as bedding material.....	187-190	relation of fungi to.....	7-8
Sheep, susceptibility to coccidioid granuloma.....	536, 538-539	seed transmission.....	55
Shive, J. W., and Martin W. H. (paper): A Comparative Study of Salt Requirements for Young and for Mature Buckwheat Plants in Solution Cultures.....	151-175	soil transmission.....	53-54
Silage, acidity.....	395-409	summering of.....	46-50
Silver maple. See <i>Acer saccharinum</i> .		Spinach. See <i>Spinacea oleracea</i> .	
Smith, L. B., and McClintock, J. A. (paper): True Nature of Spinach-Blight and Relation of Insects to Its Transmission.....	1-60	Spinach-blight. See <i>Spinacea oleracea</i> , blight of.	
Sodium hydroxide, influence on nitrogen-assimilating bacteria.....	313-332	Stahl, C. F., and Carsner, E. (paper): Obtaining Beet Leafhoppers Nonvirulent as to Curly-Top.....	393-394
Soil—		Stakman, E. C., Parker, J. H., and Piemeisel, F. J. (paper): Can Biologic Forms of Stemrust on Wheat Change Rapidly Enough to Interfere with Breeding for Rust Resistance?.....	111-124
acidity, influence on growth of legume bacteria.....	319	Stem Lesions Caused by Excessive Heat (paper).....	595-604
alkalinity, influence on growth of legume bacteria.....	319	Stem lesions of <i>Solanum tuberosum</i> —	
availability of potash in.....	297-316	causal organisms of.....	213-219
deep tilling, effect in Great Plains.....	505-517	inoculations with fungi from.....	214-219
dynamiting, effect in Great Plains.....	505-517	isolation of fungi from.....	213-214
effect of gypsum on solubility of potash in.....	65	Stemrust. See <i>Puccinia graminis tritici</i> . See also <i>Puccinia graminis tritici-compacti</i> .	
loess, moisture content of—		Straw—	
relation to hygroscopic coefficient.....	453-480	barley, use as bedding material.....	187-190
relation of weather to.....	454-457	oat, used as bedding material.....	187-190
		wheat, use as bedding material.....	187-190
		Strawberry guava. See <i>Psidium cattleianum</i> .	
		Subsoiling, Deep Tilling, and Soil Dynamiting in the Great Plains (paper).....	481-521
		Subsoiling, effect on crops in Great Plains.....	481-521
		Sudan-grass hay—	
		composition.....	176-185
		digestibility.....	176-185
		yield.....	176
		See also <i>Holcus halepensis sudanensis</i> .	
		Sugar beet. See <i>Beta vulgaris</i> .	
		Sulphuric acid. See Acid, sulphuric.	

- Sunflower. See *Helianthus annuus*. Page
- Susceptibility and Resistance to Citrus-canker of the Wild Relatives, Citrus Fruits, and Hybrids of the Genus *Citrus* (paper)..... 337-358
- Sweet potato. See *Ipomoea batatas*.
- Swine—
- Susceptibility to coccidoidal granuloma..... 536-539
- toxicity of cottonseed products to..... 449
- Synanthodon pictipes*, female, at light traps..... 146-148
- Tangelo, resistance to Citrus-canker..... 351
- Terminalia—
- calappa*, food plant of *Ceratitis capitata*..... 606-608
- chebula*, food plant of *Ceratitis capitata*..... 606-608
- Tetramesa—
- palustre*, tissue invasion by..... 569
- parvula*, tissue invasion by..... 569
- triplochis*, tissue invasion by..... 569-570
- Tetrastichus giffardianus*, parasite of *Ceratitis capitata*..... 606-610
- Thecia nerifolia*, food plant of *Ceratitis capitata*..... 606-608
- Tilling, deep, effect on crops in Great Plains..... 505-517
- Tissue Invasion by *Plasmiodiophora brassicae* (paper)..... 541-572
- Toxic culture filtrate, value in immunization against blackleg..... 261-267
- Toxicity, cottonseed products..... 435-452
- Tonju, symptomatic anthrax, concentration..... 265-266
- Toxoptera graminum*, life history..... 97-110
- Triphloxia*—
- host of *Tetramesa palustre*..... 569
- palustre*, host of *Tetramesa palustre*..... 569
- Triphasia trifolia*, resistance to Citrus-canker..... 341
- Triticum*—
- aestivum*—
- effect of—
- deep tilling in Great Plains on..... 505-517
- soil dynamiting in Great Plains on..... 505-517
- subsoiling in Great Plains on..... 484-504
- host plant of *Puccinia graminis tritici*..... 111-124
- resistance of seeds to desiccation..... 527-534
- silage of, acidity..... 400-408
- wheat-scab on, when grown following infected corn..... 611-612
- dicoccum*, host plant of *Puccinia graminis tritici*..... 118-119
- spp.—
- coefficients of correlations between—
- plant characters..... 380-389
- weight of seed and resultant plant characters..... 372-376
- effect of environment during growth..... 366-376
- seed, relation of weight to resultant plant characters..... 376-380
- variations..... 359-391
- vulgare*—
- host plant of *Puccinia graminis tritici*..... 117-121
- syn. *Triticum aestivum*.
- True Nature of Spinach-Blight and the Relation of Insects to Its Transmission (paper)..... 1-60
- Trypanosoma equiperdum*, recovery from blood of rats..... 574-576
- Trypanosoma*, recovery from blood of rats..... 571-576
- Turner, W. B. (paper): Female Lepidoptera at Light Traps..... 135-140
- Turnip, curly. See *Brassica rapa*. Page
- Two Important Introduced Parasites of the Brown-Tail Moth (paper)..... 191-200
- Vaccine, "germ-free," value in immunization against blackleg..... 253-262
- Variation and Correlation in Wheat, with Special Reference to Weight of Seed Planted (paper)..... 392-392
- Variations in the Moisture Content of the Surface Foot of a Loess Soil as Related to the Hygroscopic Coefficient (paper)..... 454-480
- Vigna sinensis*, whitespot disease of..... 598
- Waianae Mountains, coffee collection in, in 1917..... 610
- Wampa. See *Clausena wampa*.
- Water. See Moisture.
- Watercress. See *Rorippa* sp.
- Water-Holding Capacities of Bedding Materials for Live Stock, Amounts Required to Bed Animals, and Amounts of Manure Saved by Their Use (paper)..... 185-190
- Weather, relation to moisture content of loess soil..... 454-457
- West Indian medlar. See *Mimusops elengi*.
- Wheat. See *Triticum* spp.
- Wheat-scab on wheat following infected corn..... 611-612
- Whisenand, J. W. (paper): Water-Holding Capacities of Bedding Materials for Live Stock, Amounts Required to Bed Animals, and Amounts of Manure Saved by Their Use..... 185-190
- White sapote. See *Casimiroa edulis*.
- Whitespot—
- control of..... 608
- disease of conifers and other plants..... 595-604
- resemblance to damping-off disease of plants..... 595
- Wild rice. See *Zizania palustris*.
- Willard, H. F., and Pemberton, C. F. (paper): Work and Parasitism of the Mediterranean Fruit Fly in Hawaii during 1917..... 605-610
- Willow. See *Salix* spp.
- Withers, W. A., and Caruth, F. E. (paper): Comparative Toxicity of Cottonseed Products..... 545-572
- Wood, yellow. See *Ochroma elliptica*.
- Wood, influence of humidity upon..... 285-295
- Work and Parasitism of the Mediterranean Fruit Fly in Hawaii during 1917 (paper)..... 605-610
- Xanthoxylum Buxpei*, resistance to Citrus-canker..... 340
- Yellow oleander. See *Thecia nerifolia*.
- Yellow-wood. See *Ochroma elliptica*.
- Zea mays*—
- effect of—
- deep tilling in Great Plains..... 505-517
- soil dynamiting in Great Plains..... 507-508
- subsoiling in Great Plains..... 484-504
- root-rot of..... 611-612
- silage of, acidity..... 400-409
- Zizania palustris*, susceptibility of seeds to desiccation..... 526
- Zygonychus* sp., causal organism of stem lesions of *Sideman tuberosum*..... 216-219









